Formative Research for Assessing Comprehensive Primary Health Care in Mysuru City: Quantitative Survey report

<u>Supported by:</u> Health Systems Transformation Platform

<u>Organisations</u> St. John's Research Institute, Bangalore Karnataka Health Promotion Trust, Bangalore

Report by:

St. John's Research Institute, Bangalore

<u>Date</u>: March 2023



List of Contributors

St. John's Research Institute (SJRI)

- 1. Dr Prem K Mony, Prof and HOD, Division of Epidemiology, Biostatistics & Population Health, SJRI
- 2. Dr Maryann Washington, Adjunct Prof, Division of Epidemiology, Biostatistics & Population Health, SJRI
- 3. Dr Tinku Thomas, Prof and HOD, Department of Biostatistics, St John's Medical College
- 4. Dr Sumithra Selvam, Senior Resident, Division of Epidemiology, Biostatistics & Population Health, SJRI
- 5. Dr Abhilasha Nair, Assistant Prof, Division of Epidemiology, Biostatistics & Population Health, SJRI

Karnataka Health Promotion Trust (KHPT)

- 1. Mr Vidyacharan Malve, Monitoring and Evaluation Manager, Comprehensive Primary Health Care, KHPT
- 2. Mr Arin Kar, Lead, Monitoring and Evaluation, KHPT
- 3. Dr Swaroop N, Thematic Lead, Comprehensive Primary Health Care, KHPT

Health Systems Transformation Platform (HSTP)

- 1. Dr Sudha Chandrashekhar, Advisor, HSTP
- 2. Dr Kumaravel Ilangovan, Specialist Primary healthcare and PMJAY linkages, HSTP

Acknowledgement

Health Systems Transformation Platform (HSTP)

- 1. IAS Rajeev Sadanandan, Chief Executive Officer, HSTP
- 2. Dr N Devadasan, Senior Technical Advisor, HSTP

Government of Karnataka

- 1. Dr Prabhudev B Gowda, State Nodal Officer, NUHM, Department of HFW, Govt of Karnataka
- 2. Sri G Lakshmikanth Reddy, IAS, Mysuru City Corporation (MCC) Commissioner, Mysuru
- 3. Dr Nagaraju, Community Health Officer, MCC, Mysuru
- 4. Dr Prasad K.H, District Health Officer, Mysuru

List of Contents

S No.	Content	Page No
1	List of tables	3-4
2	List of figures	5-6
3	Abbreviations	7
4	Executive Summary	8-17
5	Objectives	18
6	Materials and Methods	18-26
7	Results	27-70
8	Out of Pocket Expenditure results	71-86
9	Conclusions	87-91
10	ICD Mortality Data	92-95
11	Appendices (1-8) (Facility and community	96-140
	questionnaires)	

List of Tables

Table No.	Title of the table	Page No
1	Number of facility and field health workers surveyed from public	21
	and private health facilities.	
2	Randomly selected wards in Strata 1 and strata 2	23
3	Details of tools used for data collection	25
4	Characteristics of health facilities	27
5	Infrastructure and equipment available in health facilities	28
6	Availability of health information technology at public facilities	29
7	Staff welfare activities in public facilities	30
8	Maternity and child health services available at facilities	31
9	Reported adult health services of public and private facilities	32
10	Adult emergency and referral services reported by public and private facilities	33
11	Monthly load of services of public facilities based on record	33
11	review of the last month	33
12	Lab investigations performed in a month based on record review	34
	of the last month	
13	Median number of persons reached for health services at the	35
	community by UPHCs	
14	Health workers filled against required Standards for UPHCs	36
15	Demographic variables of health workers from UPHCs	37
16	Training received as reported by doctors and nurses from public facilities	37
17	Training reported to have been received by lab technicians and	38
10	pharmacies at all public facilities	20
18	Training reported to have been received by ANMs and ASHAs in public facilities	39
19	Role of doctors at public facilities	40
20	Role of nurses at public facilities	40
21	Demographic characteristics of the study population	41
22	Descriptive statistics of age group by gender of the population	42
23	Distribution of age group <5 years by gender of the population	42
24	Demographic parameters of the surveyed population	42
25	Household Characteristics of the surveyed population	43
26	Health insurance statistics of the surveyed population	44
27	Lifestyle characteristics of the surveyed population	45
28.1	Health seeking behaviour characteristics in 2 weeks morbidity	47
28.2	2 weeks morbidity – Distance, time and cost involved	48
29.1	Maternal health: Health seeking behaviour characteristics	50
29.2	Maternal health: Cost of care and satisfaction score	51
30	Child health: Health seeking behaviour, and satisfaction score	52
31	Health care seeking characteristics for child immunization	52
32.1	Non-Communicable Diseases: Health seeking behaviour characteristics	53
32.2	Non-Communicable Diseases: Details of investigations and complications	54

32.3	Non-Communicable Diseases: Recent investigations and measurements details	55
32.4	Non-Communicable Diseases: Managing routine medical expenses and satisfaction scores	55
33.1	Comparison of demographic parameters between individuals having pucca and non pucca houses	56
33.2	Comparison of 2-week morbidity characteristics between individuals having pucca and non pucca houses	57
33.3	Comparison of maternal health characteristics between individuals having pucca and non pucca houses	58
34.1	Comparison of demographic parameters between households belonging to wards with predominant notified slums and non-slum areas	58
34.2	Comparison of 2-week morbidity characteristics between individuals belonging to wards with predominant notified slums and non-slum areas	59
34.3	Comparison of maternal health characteristics between individuals belonging to wards with predominant notified slums and non-slum areas	59
35.1	Comparison of health insurance between households showing and not showing BPL card at the time of survey	60
35.2	Comparison of 2-week morbidity characteristics households showing and not showing BPL card at the time of survey	60
35.3	Comparison of maternal health characteristics between individuals showing and not showing BPL card at the time of survey	61
36	Socio-demographic characteristics of patients from public and private facilities	61
37	Overall satisfaction on services as reported by patients at exit interview	65
38	Association between the Reimbursed and Non- reimbursed categories with various Socio demographics characteristics	80
39	Association of type of Medical Institution and Mean Expenditure	85
40	Association of Place of Residence and Mean Expenditure	85
41	Association of Various Components of Medical Expenditure and Type of Medical Institution	85
42	Association of Wealth Quintiles and Mean of components contributing to Medical Expenditure - Inpatients	86
43	Association of Wealth Quintiles and Mean of components contributing to Medical Expenditure – Outpatients	86
44	ICD Classification by gender	92
45	ICD Classification by age category	94

List of Figures

Figure No.	Title of the figure	Page No
1	Map showing location of UPHCs in Mysuru city.	19
2	Selection of private health facilities	20
3	Process of sampling method for community survey	22
4	Mysuru city map- highlighting the wards studied	23
5	Reported outreach services provided by public facilities	35
6	Types of ailments in the surveyed population in the last 2 weeks	46
7	Place of health care for different ailments in the surveyed population in the last 2 weeks	46
8	Distribution of individuals who sought treatment at health facility by age group	48
9	Distribution of individuals according to type of facility from where they received treatment	48
10	Patients exit interview: Health problem for the current visit to the facility	63
11	Patients exit interview: Reason for choice of facility for current visit	63
12	Patients exit interview: What is liked most about the health facility by patients	64
13	Patients exit interview: Services received at the facility for current visit	64
14	Patients exit interview: Level of satisfaction for services received at the health facility	65
15	Challenges faces by HCPs from UPHCs / doctors of private facilities	66
16	Challenges faced by field HCPs from UPHCs	67
17	Suggestions given by HCPs from UPHCs / Doctors from private health facilities	68
18	Suggestions given by field HCPs from UPHCs	69
19	Suggestions given by patients at exit interviews for improving services at public and private facilities	70
20	Distribution of Non-Communicable Diseases among female inpatients	72
21	Distribution of Non-Communicable Diseases among male inpatients	72
22	Distribution of Non-Communicable Diseases among female outpatients	73
23	Distribution of Non-Communicable Diseases among male outpatients	73
24	Source of total expenditure for inpatient admissions	74
25	Source of total expenditure for outpatient care	74
26	Expenditure component for Inpatients	75
27	Expenditure component for Outpatients	75
28	Expenditure breakup for Inpatients	76
29	Expenditure breakup for Outpatients	76
30	Details of medical expenses in Inpatients	76
31	Details of medical expenses in Outpatients	76

Formative Research for CPHC in Mysuru City

32	Distribution of inpatient medical expenditure	77
33	Components of medical expenditure for Inpatients by wealth quintiles	78
34	Components of medical expenditure for Outpatients by wealth quintiles	78
35	Distribution of NCDs reported by age among Inpatients	79
36	Distribution of NCDs reported by age among Outpatients	79

Abbreviations

ADD	: Acute Diarrheal Diseases
AFB	: Acid Fast Bacilli
ANC	: Ante Natal Care
ANM	: Auxiliary Nurse Midwife
ARI	: Acute Respiratory Infections
ASHA	: Accredited Social Health Activist
BP	: Blood Pressure
CEB	: Census Enumeration Block
CHC	: Community Health Centre
COPD	: Chronic Obstructive Pulmonary Disease
CPHC	: Comprehensive Primary Health Care
CGHS	: Central Government Health Scheme
C-section	: Caesarean section
DKA	: Diabetic Keto Acidosis
ESI	: Employee State Insurance scheme
eVIN	: Electronic Vaccine Intelligence Network
HCPs	: Health Care Providers that included those in the facility (doctors, nurses, lab
	technicians and pharmacist and those in the field that included (ANMs and ASHAs)
KSMSC	: Karnataka State Medical Supplies Company
MCH	: Maternal and Child Health
NCD	: Non-Communicable Disease
NVBDCP	: National Vector Borne Disease Control Program.
OOPE	: Out-of-Pocket Expenditure
PHC	: Primary Health Centre
PNC	: Post Natal Care
TB	: Tuberculosis
UID	: Unique Identification Data
UPHC	: Urban Primary Health Centre

Executive Summary

The overall aim of the Formative Research was to assess the status of urban Comprehensive Primary Health Care (CPHC) in Mysuru city. This part of the pre-final report covering the quantitative survey by St John's Research Institute, Bangalore, addresses the following specific objectives:

- 1. To identify and explore the role of key stakeholders in the provision of urban primary health care.
- 2. To describe the status of urban comprehensive primary healthcare (UPHC) in Mysuru city at three different levels health systems, facility, and community
 - a) at health system level for capacity for designing, developing, implementing, and monitoring urban primary health care in Mysuru city.
 - b) public and private health facility readiness for delivering preventive and nondomiciliary curative primary health care in urban Mysuru.
 - c) profile the community morbidity status, healthcare seeking, and costs incurred for selected acute and chronic conditions in urban wards of Mysuru city.
- 3. To identify and explain barriers and facilitators to comprehensive Primary Health Care

Methodology of the Community Assessment:

The community assessment was carried out in 25 wards of Mysuru city. Study design was population-based survey representative of Mysore city. Sampling design was as follows: The total number of wards surveyed in urban Mysuru was 25 out of 65 wards. The wards were divided into two strata, the first strata (STRATA1) consisted of wards with 5 to 20 CEBs each and the second strata (STRATA2) contained wards with more than 20 CEBs each. Out of 32 wards in STRATA1, 12 wards were randomly selected. Similarly, 13 wards were randomly selected from STRATA2 (total 33 wards). From the selected wards, 192 and 208 census enumeration blocks were randomly selected (16 CEBs in each ward). At the final stage, 15 households per CEB were selected by systematic sampling. The total sample size was 6000 households to conduct the community survey.

Methodology of the Health Facility Assessment:

All the public primary health facilities (20 UPHCs) and 20 private health facilities along with all the three public health and three private health facilities (<30 bedded) offering childbirth services and functioning 24/7 from Mysuru city. Sampling method was as follows: Health Care providers (HCP) - one doctor, nurse, lab technician, pharmacist,

ANM and ASHA available at each facility selected purposively based on availability and seniority. Patients – four from each facility selected purposively. Data collection was from May to Sept 2022. Survey with HCPs to assess their roles, challenges, and suggestions for improving CPHC, record review to assess reach of services, observation checklist for amenities, equipment, supplies at the health facilities.

Results of the Community Assessment:

A total of 6007 households comprising of 21576 individuals were surveyed from 25 randomly selected wards of Mysuru city. Out of total population surveyed, 89.2% belonged to less than 60 years of age and 55.5% of them were above 30 years of age. Equal proportion of males (48.8%) and females (51.2%) were there in the surveyed population. One third of the population had education up to middle school level, and 1.1% of them were either widowed/ separated / divorced. Around 40% of them were employed. Among 6474 women in the reproductive age group, 100 (1.6%) mothers were currently pregnant and pregnancy in the last 3 years was 563 (8.6%). Among 11978 individuals aged above 30 years, 17.3% of were either diabetes or hypertension. Approximately 82% of them had pucca house, 99% had improved toilet facility and 93% had improved water source which was comparable to NFHS-5 data.

Health Insurance: Two third of the households (67.8%) didn't have any insurance coverage. Only 17% of the households were covered under Ayushman Bharat / Aarogya Karnataka. ESI / CGHS and private insurance coverage was 7.8% and 7.4% respectively. Only 26 households utilized their health insurance in the past one year.

Lifestyle of adults: Tobacco consumption in both smoking and chewing form was observed in 3% of the population respectively among individuals aged above 18 years. Alcohol consumption was reported in 4.4% of the individual's majority being males. Only 5% of the individual noted to be part of some voluntary organization.

2 weeks morbidity status: Illness in the last 2 weeks was observed in 7% (n=1490) of the population studied, of which almost half of them sought treatment at health facilities indicating moderate utilization of health facilities for acute illnesses. Among those who sought treatment, only 32.5% received treatment from public health facility. Self-

medication and use of Over the Counter (OTC) drugs were the reasons reported among those who did not seek treatment at health facility.

Less facilities and long distance were the reasons reported for changing place of treatment among 9% of the people who sought treatment at multiple health facilities. No significant difference between public and private facilities in terms of change of place of treatment was observed. Distance to health facility, time taken to reach the health facility and time taken to consult the doctor were comparable between individuals seeking care at public and private health facilities, indicating the preference for choosing health facility was not governed by the above said factors. Income and savings were the most used mode for managing their routine medical expenses and were comparable between individuals choosing public and private health facilities. The preference for health facilities was comparable by gender, however, significant difference was seen by age categories. Significantly higher proportion of children between 6-18 years were consulted in private health facilities, which could be because of availability of paediatric specialist in the private set up.

Among the people who utilized PHCs, District Hospital and ESI hospital, higher proportion (~60%) belonged to middle aged and elderly. Families preferred private practitioners, clinics, and hospitals (~30%) for the ailments in their children and adolescents. Preference of health care facilities for their ailments in the past 2 weeks showed that higher proportion of people with morbidity of musculoskeletal pain, respiratory problems and for general weakness, preferred public health facilities for health care. For Non-Communicable Diseases, private hospitals were most preferred health facility.

Approximately 60% of people residing in the non-slum area preferred private clinics and hospitals for their ailments. Urban PHCs were the next place of preference for non-slum population. Utilization of district hospital was noted be higher among population belonging to slum area. Even in slum population, 50% of them preferred private health facilities.

Maternal health: Antenatal care (ANC) was elicited only among the current pregnancies (n = 100) during the study period. All these pregnancies were registered and 46% of them had reported utilizing public health facilities for ANC care. The data on childbirth was recorded from the mothers who delivered in the past 3 years. More than half of the mothers

utilized public health facilities for their deliveries, and a 53% of them had normal vaginal Childbirth. Significantly higher proportion of females had C-section in private health facilities (67%) compared to only 25% in public health facilities (p<0.01). Similarly for Postnatal care also, 55% of them preferred public health facilities.

The reasons for choosing the public health services for maternal health care were reported to be near distance and free of cost. Good doctor, timely service, and all facilities available at one place were the primary reasons for preferring private health facilities. Like the pattern observed for acute illnesses, the distance, time, and cost spent were not determining factors for choosing MCH facilities.

Child Health (\leq 60 months): About a quarter of the children (25%) were reported to be sick in the last 1 month. Acute Diarrhoeal Disease (ADD) (67.6%) was the most reported illness followed by Acute Respiratory Infection (ARI) (35.5%). For both the ailments, approximately equal proportion of households availed care from public and private health facilities. The preferred reasons for choosing public health facility were less / free of cost followed by trust in doctor and nearby distance which was similar for both ARI and ADD. Trust in doctor, timely service, and all facilities at one place were the reasons reported in favour of private health facility. Hospitalization rate for ARI and ADD were 13.2% and 4.0% respectively. Majority of them were hospitalized in private health facilities (ARI – 80% and ADD – 70%). Due to free cost of immunization, majority of children (< 2 years) have been reported to avail child immunization services in the public health facilities. These findings indicate that although the preference of health facilities for outpatient care was equal in both public and private, but for hospitalization, majority of them preferred private health facilities. This could be due to the availability of comprehensive paediatric care in a private setting.

Non-Communicable Diseases (>30 years): The reported prevalence of either diagnosed diabetes or hypertension was 17.3% (15.9% in males / 18.7% in females). Both diabetes and hypertension were presented in 7.5% of the individuals (6.5% males/ 8.7% females). Diabetes and Hypertension alone was reported in 12.8% and 15.0% respectively.

For NCD care, private health facilities were the preferred health facilities, considering the trust in doctor (80.0%) followed by timely service (50.0%) and all facility at one place (27.0%). Higher proportion of both diabetes (70.3%) and hypertension (65.9%) patients

preferred private facility for buying medicine routinely. Even for the NCD complications, most of them were referred to private health facilities.

Non availability of NCD drugs round the year and lack of investigation facilities may be implicated as the reason for inclination towards private health facilities. Like other illnesses, income and savings were reported to be the commonest mode of managing routine medical expenses in NCD patients.

Health economics: The cost incurred for healthcare in public health facilities was very less as compared to private health facilities. Although there was no / minimal charge of consultation in public health facilities, the median investigation cost and the drug cost was Rs. 65 (10, 520) and Rs. 110 (0, 425) respectively. The distribution of Out-of-Pocket Expenditure (OOPE) per person / ailment was \leq Rs 500 (57%), 500-1000 (21.6%), 1000-5000 (15.9%) and > 5000 (5.4%). Considering the family income availability, catastrophic OOPE (>10% of the annual income) was observed in 8 of the surveyed households.

Regarding the cost spent towards ANC care, those who preferred private health facilities had reported to spend five folds of what was spent in public health facilities (Median cost in Pvt = Rs 15,000, Govt. = Rs 3,000). For all maternal health services, income and savings were the most common utilized modes for managing medical expenses. Only 1% of them utilized health insurance for their childbirth purpose. The cost spent towards childbirth care was significantly higher among those who utilized private health facilities compared to public health facilities (Median cost in Pvt = Rs 50,000, Govt. = Rs 5,000). PNC care expenses were also noted to be higher in the private as compared to public health facilities. The median cost spent on treatment for both ARI and ADD in public was one third of what was spent in private health facility.

Satisfaction Score: Satisfaction score was assessed for NCDs and MCH services. People who had utilized public and private health facilities reported median score of 8 and 9 respectively.

Analysis was done to compare health seeking behaviour between wards with and without predominantly slum areas. The pattern of health seeking was similar in both slum and non-slum population. While comparing population who had shown and not shown BPL card during survey, higher proportion of public health facilities were utilized by people who showed BPL card.

Results of the Health Facility Assessment:

Characteristics of health facilities: Median population covered by UPHCs was 50097 and slum population was 6407. UPHCs were functional for 6 hours during daytime while most of the private clinics (95%) were primarily functional in evening hours. Nearly 70% of UPHCs were branded as Health and Wellness Centres, 30% of private clinics offer integrated medicine. Public health facilities were on an average distance of 1km from patients' residence while private health facilities were on an average of 2km distance. The patients from public and private differed significantly (p<0.05) by sociodemographic characteristics such as age (younger in private facilities); sex (more males seeking services at private facilities); occupation (lesser homemakers among those seeking services from private facilities) but not by education level. The commonest health problem for current visit of patients to the health facility was fever (30% and 39%) in both public and private health facilities; pain (25% in both). Few patients returned for follow-up or check-up of diabetes (11% and 16%) and hypertension (11% and 10%) from public and private health facilities. The commonest reason for choosing public health facility was free treatment (29%) and good response of health workers (39%), while for private health facilities it was good consultation (54%) and nearby location (27%).

Availability of amenities, essential drugs, and equipment: Waiting area was available in 100% of public and 80% of private health facilities. Toilets was available in all public UPHCs, public and private childbirth facilities. There was limited availability of space within and beyond (50-meter radius) all the health facilities for yoga practice, counselling, and nutrition demonstration. More than 70% of all facilities had pharmacies; >56% had a clinic and >40% had a lab on the same street. Basic CPHC equipment (BP apparatus, glucometer, weighing machine, pulse oximeter) were available in public and private facilities. While ophthalmoscope, Snellen's chart, cardiopulmonary monitor were not available in 95% of UPHCs. Essentials drugs to manage minor health issues and treatment of NCDs such as diabetes and hypertension were available in all public health facilities. Essential drugs for initial management of obstetric and cardiac emergencies, prevention of cardiac and neurological complications was not available in >60% of public health facilities. Less than 25% of public and none of private health facilities had a UID for registration of patients, despite computer and online support being available in 100% of

public health facilities. Around 25% of UPHCs provided all staff welfare activities such as immunisation, post exposure prophylaxis and annual health checks.

Services provided by health facilities: All UPHCs and 35% of private clinics offered maternity (antenatal) services. More than half of the UPHC's and 10% of private clinics offered neonatal services. More than 95% of public and only 15% of private clinics offered child health services. Less than 35% of UPCHs provided services for cancer, mental health and endocrine issues. More than half of the private clinics offered services for NCDs-diabetes and hypertension.

Load of services provided monthly by health facilities: Median monthly OPD registration of UPHCs was 1447 and of childbirth public and private health facilities was 1952 and 1140 respectively. Median monthly ANC registration was 26 in UPHCs and 353 in childbirth public but only 17 in private childbirth facilities. Two outreach services were conducted by UPHCs. Median patients treated at UPHCs, public and private childbirth health facilities for diabetes (99, 74, 160), hypertension (138, 70, 160) and TB (14, 23, 23) respectively. Median pregnancy tests at public UPHCs, public and private childbirth facilities was 10, 40 and 10 respectively; 169, 709 and 51 random blood sugar tests were performed respectively. Lipid profile, thyroid test, Renal test, and dengue test were performed only by private childbirth facilities. ANMs reported a current average of 40, 171 and 180 pregnant women, diabetics or hypertensive patients being followed up while ASHA reported a current average of 7, 211, 212 respectively.

Status of HCPs at public health facilities: Except for doctors and lab technicians, there was shortage of 27% nurses, 17% of ANMs, 80% of ASHAs, 25% of pharmacists, 60% of DEOs and 35% of Class D workers. More than 90% of nurses, lab technicians and ASHAs were employed on contract basis. All 100% of nurses, ANMs and ASHAs, 59% of lab technicians and 88% of pharmacists were females. Amongst doctors and nurses, 35% and 52% respectively received training on SBA; 48% and 30% respectively received training on RBSK and 52% and 35% respectively received training on RKSK over the last 5 years. While only 35% and 33% of ANMs received this training on RBSK and RKSK respectively. More than three quarters (78%) of lab technician received training on NVBDCP and 57% on TB over the last 5 years, 9% mentioned they had stock out of lab supplies in the last three months. More than half (55%) of pharmacists received training on eVIN over the last 5 years, 44% mentioned they had medication stock outs.

Perceptions of patients on services received: The satisfaction score of private childbirth facilities (73.9 ± 11.1) was higher than that of public facilities (69.2 ± 11.8) , but this was not significantly different. However, patients from UPHCs (71.3 ± 10.7) were significantly more satisfied with services received than those from private clinics (58.3 ± 17.2) at p<0.0001.

Challenges faced by HCPs:

Health system related challenges -- HCPs of health facilities reported the following challenges: Less ratio of HCP with population (35%), multi-tasking job (20%), lack of supplies and meeting targets (19%), clinical management (37%) by HCPs of private facilities. Challenges reported by field level HCPs were management of targets (28%), ratio of HCP and population (23%) and multi-tasking (21%).

Community related challenges -- By HCPs of health facilities: lack of cooperation of people (44%). Lack of acceptance of people to treatment protocols (41%). By field level HCPs: lack of cooperation of community (41%) and difficulty mobilizing the community (21%).

Suggestions to improve services:

Health system related suggestions mentioned by edHCPs from UPHCs and private facilities: By HCPs of public and private health facilities: Building improvement (63%), drug availability (34%), better lab facilities (30%), better equipment and facilities (23%). By field level HCPs: Reimbursement (46%), better ratio of HCPs with population (38%) and better salary (35%). By patients of public health facilities: lab equipment improvement (18%), need for specialists (9%), inpatient facility for emergencies (8%). By patients of private health facilities: lab equipment (4%), need for specialists (5%), inpatient facility for emergencies (2%).

*Community related suggestions mentioned by HCPs--*By HCPs of health facilities - On the spot treatment as well as better hygiene by 8%; 35% mentioned health education for the community.

By field level HCPs >27% - mobilization of people through key stakeholder involvement.

Secondary data analysis of National Sample Survey data on Out-of-Pocket Health Expenditure in Karnataka:

Using NSSO data (Round 75, 2017-18), health expenditure towards NCDs in Karnataka was analyzed. Cost of medicines was the major expenditure for both inpatients and

outpatients. The association of reimbursement categories (reimbursed or not reimbursed) with various socio demographic characteristics was analyzed. Social group, education, type of health facility, place of residence, wealth quintiles and employment category had significant association with reimbursement. Individuals belonging to SC, ST and OBC categories, being literate, individuals preferring private hospitals, urban residents, individuals belonging to highest wealth quintile and self-employed, non-agriculturists had proportionately higher reimbursement. Average expenditure of consultation, drugs and investigation cost were significantly higher for the private health facilities. Average expenditure for inpatients was higher for individuals belonging to urban and higher wealth quintiles.

In summary, health facilities were easily accessible to the population. Regular supervision and monitoring of HCPs by a senior within the health facility or health office was occurring. Basic services of ANC, management of minor ailments, first aid for injuries was being managed by UPHCs and private clinics despite HCP shortage. Patients were satisfied with services received and accessed services based on proximity and their perception of HCPs. Leadership and governance need to focus towards improving quality of care rather than just quantity. The availability of services was limited to 7 hours by all the UPHCs and 5 hours by the private clinics. Only those facilities that provided childbirth services were functional 24/7. Both facility and field HCPs highlighted the need to improve the building / equipment / lab / maternal services. Services at public health facilities were mostly accessed by homemakers and women while the private health facilities were accessed mostly by males and younger age group. More robust health information system that not only facilitates registration of patients so that follow-ups and linkages between facility and field HCWs are planned strategically especially for those with chronic NCDs and CDs but would also aid in monitoring progress with meeting targets. Feedback from the community or individuals could be obtained to determine ways to improve access, quality, and availability of services. Capacity building of HCWs at all levels must be geared towards better communication with patients, identification of complications and appropriate referrals, linkages between public and private health facilities that probably use a common UID for patients to facilitate efficient follow-ups. Capacity building could be facilitated by using the mentoring approach. It would be prudent for public health facilities to be re-organised so that there is at least one facility offering childbirth services attached to 4-5 UPHCs. Moreover, given the health workforce shortage, a system to make diagnostic services more efficient, yet accessible could include sample collection at the UPHCs with an effort to transport samples to a referral diagnostic centre that would report back to the UPHC details of the test result. This will require a better health information system that links all UPHCs with the referral diagnostic centre.

Main Report

A. Objectives of the assessment of primary health care services in Mysuru city

Overall objective of formative research was to describe the status of urban comprehensive primary health care system in Mysuru city, identify and analyse barriers and facilitators to comprehensive primary health care, and identify design options to strengthen urban primary health care.

A.1 Specific Objectives

- 1. To identify and explore the role of key stakeholders in the provision of urban primary health care.
- 2. To describe the status of urban comprehensive primary healthcare (UPHC) in Mysuru city at three different levels health systems, facility, and community
 - a) at health system level for capacity for designing, developing, implementing, and monitoring urban primary health care in Mysuru city.
 - b) public and private health facility readiness for delivering preventive and nondomiciliary curative primary health care in urban Mysuru.
 - c) profile the community morbidity status, healthcare seeking, and costs incurred for selected acute and chronic conditions in urban wards of Mysuru city

To identify and explain barriers and facilitators to comprehensive Primary Health Care and
 To identify design options for strengthening urban primary health care.

B. Methods:

As part of a 5-month formative technical support project (April-August 2022), to assist the Government of Karnataka to strengthen comprehensive urban primary health care, Mysuru city corporation was selected. Mysuru city has a total population of 1261,000 as of 2022, with a 2% 1993 plateaued approximate increase in population since (https://www.macrotrends.net/cities/21343/Mysuru/population). We carried out a quantitative study with a [i] facility assessment and service availability survey of all urban public health facilities and similar private health facilities in Mysuru city corporation. [ii] community assessment through a general household survey, to assess profile of community morbidity status, their healthcare seeking behaviour and costs incurred for selected acute and chronic conditions in urban wards of Mysuru.

B.1 Health facility assessment

<u>B.1.1 Sample:</u>

Selection of public health facilities: All 23 public urban primary health centers were selected for the assessment. It consisted of 20 UPHCs and three centers offering childbirth services.



Figure 1. Map showing location of UPHCs in Mysuru city.

Selection of private health facilities: An equivalent number of private health facilities were selected. Initially the list of health facilities commonly utilized by the community that were elicited from the respondents in the community survey showed that the top health facilities mentioned were either public or private secondary or tertiary level hospitals (Figure 2). Hence from the rest of health facilities named (n=174), eight were not eligible to be included since they had >35 beds. Thus, of a total of 168 health facilities, 85% were excluded for the reasons mentioned (Figure 2). Twenty-three health facilities – 3 hospitals offering childbirth services with less than 35 in-patient beds and 20 clinics were selected purposively if they consented to participate in the study.




Figure 2: Selection of private health facilities

Selection of health care providers (HCP): From each public health facility, one doctor, nurse, lab technician, pharmacist, ANM and ASHA worker were selected, based on availability and their consent to participate in the survey (Table 1). Similarly in the private health facilities, based on availability HCPs were selected to assess their roles, challenges to provide and suggestions in improving comprehensive primary health care package of services.

		I	Field HCPS			
	Doctors	Nurses	Lab technician	Pharmacist	ANM	ASHA
Clinics / UPHCs ✓ Public UPHCs (n=20) ✓ Private Clinics (n=20)	19* 20	20 0^	20 0^	20 ^x 0 [^]	20 0^	19^^ 0^
Facilities offering Childbirth services: ✓ Public (n=3) ✓ Private (n=3)	3 2	3 3+	3 3	3++ 3 ^{xx}	1 ^{xxx} 0^	1 ^{xxx} 0^

Table 1: Number of facility and field health workers surveyed from public and private health facilities.

*Kumbarakoppal- doctor did not complete;

^: No nurse, lab tech, pharmacist, ANM, ASHA available in private clinics;
⁺¹ is an ANM;
⁺⁺1 is a Diploma Nurse;
^x: 3 are Diploma Nurses and 2 are Medical Officers performing the pharmacist role too, took the survey in 5 UPHCs;

^: 1 UPHC, ASHA was not available

xxx: No ANM/ASHA for two public health facilities offering childbirth services.

Selection of patients: Four patients were selected based on availability and their consent to participate in the study.

B.2. General Household Survey

Multi-stage stratified random sampling was used to identify the sample households.

Sampling method

The universe was the citizens residing in the 65 wards of the Mysuru City Corporation. Each ward was further subdivided into Census Enrolment Blocks (CEB). Those wards with less than 5 CEBs per ward will be excluded from the study sample. The total number of wards surveyed in urban Mysuru was 25 out of 65 wards. It was a 3-stage process to obtain the required sample size of 6000 households. The wards were divided into two strata, the first strata (STRATA1) consisted of wards with 5 to 20 CEBs each and the second strata (STRATA2) contained wards with more than 20 CEBs each. Out of 32 wards in STRATA1, 12 wards were randomly selected. Similarly, 13 wards were randomly selected from STRATA2 (total 33 wards).

From the selected wards (Table 2), 192 and 208 census enumeration blocks were randomly selected (16 CEBs in each ward). At the final stage, 15 households per CEB were selected by systematic sampling total 6000 households were chosen to conduct the community survey. (Figure 3)




Figure 3: Process of sampling method for community survey

			Total no			Total no
Sno	STRATA1	Ward No	of CEB's available	STRATA2	Ward No	of CEB's available
1	WARD No0010	10	20	WARD No0001	1	30
2	WARD No0023	23	23	WARD No0003	3	26
3	WARD No0027	27	23	WARD No0007	7	37
4	WARD No0028	28	23	WARD No0011	11	26
5	WARD No0029	29	21	WARD No0020	20	28
6	WARD No0030	30	20	WARD No0021	21	27
7	WARD No0031	31	21	WARD No0026	26	25
8	WARD No0037	37	24	WARD No0033	33	25
9	WARD No0038	38	21	WARD No0040	40	27
10	WARD No0041	41	21	WARD No0049	49	29
11	WARD No0047	47	22	WARD No0054	54	25
12	WARD No0060	60	16	WARD No0055	55	26
13				WARD No0058	58	28

 Table 2: Randomly selected wards in Strata 1 and Strata 2




Figure 4: Mysuru city map- highlighting the wards studied

There were 192 CEBs in the first strata and 208 CEBs in the second strata. From each CEB, 15 households were randomly selected, giving a total of 6,000 households. These households were listed, and those with a patient with acute illness or a diabetic/hypertensive were chosen for the study.

Data analysis

Descriptive statistics were reported as number and percentages for all the categorical variables, median (IQR) for all continuous variables when not normally distributed or as mean (\pm SD) of both facility and community assessment. The cost of care analysis (for 2 week morbidity, Maternal and Child health services, Diabetes and Hypertension care) and satisfaction scores were reported as median with 25th and 75th percentiles.

B.3. Tools used for data collection

B.3.1. Tools for Facility Assessment

The tools were developed iteratively by a team of public health experts, doctors, nurses and in keeping with Standards for Urban Primary Health Care (Ministry of Health and Family Welfare, 2015). In all there were four forms to collect information relevant to meet the objectives of the study (Table 3). Two field investigators were trained by the core team to collect the information after they obtained informed consent from the head or medical officer of the health facility.

Form 1: This consisted of 8 sections as mentioned below.

- <u>Section A: Health Facility General Information</u>: This was completed by the trained field investigator and gave information on availability of CPHC services, specialists, and accessibility for patients
- <u>Section B: Health workers- Sanctioned and Filled (all available HP):</u> Total number of available health workers was elicited using this section
- <u>Surveys: Section C- Doctors; Section D Nurses; Section E Lab Technician; Section F –</u> <u>Pharmacist; Section G – ANMs; Section H – ASHAs [1 per facility]</u>: The roles, suggestions for improvement of CPHC and challenges faced by health care personnel was obtained.

Form 2: Facility Audit: This was an observation checklist which provided information on infrastructure, equipment and supplies including drugs and lab tests available at the health facility. *Form 3: Record Reviews – Services and Load of Services:* Information on services available and monthly load of these services were elicited through review of records available at the health facility and with the help of the relevant health care personnel.

Form 4: Patient Exit Interviews: Information on reasons for visiting the health facility, choice of health facility, services received, satisfaction with services received and suggestions for improving the present services.

Tools used	Purpose	What method used	From whom
Form 1:	•		
Section A:	To obtain general information of the health facility: Type of facility, location, catchment population, intersectoral coordination, services available, health worker availability with timing and finance	Interviewed	Medical officer or senior nurse
Section B	To obtain information on personnel such as number of sanctioned health care providers (HCPs), number available	Interviewed Record review	Medical officer or senior nurse
Section C	To obtain information from one doctor on experience, education, training received, supervision and monitoring, challenges faced, roles performed, confidence in performance of roles, and suggestions for improvement	Interviewed	Medical officer or available duty doctor
Section D	To obtain information from one nurse on same topics as given in Section C	Interviewed	Senior nurse or nurse on duty
Section E	To obtain information from one Lab technician on same topics as given in Section C. In addition, information on stock outs and equipment's not working	Interviewed	Lab technician available
Section F	To obtain information from one pharmacist on same topics as given in Section C. In addition, information on stock outs and medicines not dispensed	Interviewed	Pharmacist or acting pharmacist
Section G	To obtain information from one ANM on training, outreach services provided, reach of services, population served, supervision and monitoring, roles, confidence in performing roles, challenges, and suggestions to improve services	Interviewed Record review	Senior ANM or ANM available
Section H	To obtain information from one ASHA on training, outreach services provided, reach of services – number followed up, number currently registered, population served, reports / registers maintained, supervision and monitoring, roles and confidence in	Interviewed Record review	Senior ASHA or who was available

 Table 3: Details of tools used for data collection.

	performing roles, challenges faced, and finally on suggestions to improve services.		
Form 2	To obtain information on the health facility – Facility Audit	Observation by walk-through Interview	Senior nurse / pharmacist/ lab technician
Form 3	To obtain information on load of services	Record review Interview	Senior nurse/ pharmacist/lab technician/DEO
Form 4	To obtain information from patients who sought services in the health facility on purpose of seeking service, distance of health facility from home, suggestions to improve services, satisfaction on services received, services received for current visit	Interview	Any patient who received services at the health facility – 4 per health facility as available

B.3.2 Tools used for Community Assessment Tools used for community survey have been appended.

C. Results:

C.1. Public and private health facility readiness for delivering preventive and non-domiciliary curative primary health care in urban Mysuru.

This section provides information on the total population covered, functioning time and type of health facility (Table 4), describes infrastructure, equipment, and supplies of the health facilities (Table 5-6); Staff welfare facilities (Table 7); services available and provided as reported by HCPs (Table 8-13; Figure 5).

C.1.1. Characteristics of health facilities

Table 4: Characteristics of health facilities

	Public		Pr	ivate
	UPHCS (n=20)	Childbirth facility (n=3)	Clinics (n=20)	Childbirth facility (n=3)
Total Population [Median]	50097			
Slum Population [Median]	6407			
Operating in own building [No (%)]	17 (85%)	3 (100%)	5 (25%)	1 (33%)
 Functioning time ✓ 24/7 ✓ 9 hours + on call ✓ 6 hours ✓ Morning hours only ✓ Morning and evening hours ✓ Evening hours only 	20 (100%)	1 (33%) 2 (67%)	1 (5%) 10 (50%) 9 (45%)	3 (100%)
 Type of facility [No (%)] ✓ Health and wellness ✓ UPHCs ✓ Childbirth UPHC (10 beds) ✓ Childbirth hospital (avg 24 beds) ✓ Childbirth CHC (30 beds) ✓ Integrated medicine clinic ✓ Allopathy medicine clinic 	14 (70%) 6 (30%)	1 (33%) 2 (67%)	6 (30%) 14 (70%)	3 (100%)

Most public facilities were operating in their own building.

Only public UPHCs reported on the median population (10600 and 3121) and slum population (1500 and 1900) as covered by ANMs and ASHAs, respectively.

Equipment such as BP apparatus, glucometer, thermometer, weighing machine, pulse oximeter, to cover basic CPHC services were mostly available in all public and private facilities. However, equipment such as cardiopulmonary monitors, ECG machine to identify any emergencies or ophthalmoscope and Snellen's chart to assess eyes were available in less than 30% of the health facilities as seen in Table 5. Certain lab equipment such as biochemistry analyzer were available in 20% of UPHCs (Table 5).

All the UPHCs and public childbirth facilities had basic essential drugs to manage minor health issues and for treatment of NCDs such as diabetes and hypertension. However, 3/20 (15%) of UPHCs did not have injection dexamethasone, 1/3 (33%) of public childbirth facility did not have injection Magnesium Sulphate, 19/20 (95%) of UPHCs did not have Injection Oxytocin all of which are useful for initial management of maternal complications. Tab Clopidogrel, a drug useful to prevent heart attacks and stroke amongst persons with heart disease (recent heart attack), recent stroke or blood circulation disease was not available 18/20 (90%) and 2/3 (67%) of UPHCs and public childbirth facilities. Emergency drugs that were not available included Inj. Calcium Gluconate in 14/20 (70%) of UPHCs, Injection Adrenalin and Inj. Hydrocortisone in 1/20 (5%) of UPHCs. Tab Aspirin was not available in 13/20 (65%) and 1 (33%); while Statins were not available in 9/20 (45%) and 1/3 (33%) of UPHCs and public facilities with childbirth services respectively. Antihypertensives such as Tab Enalapril was not available in 4/20 (20%) of UPHCs and Tab Losartan was not available in 12/20 (60%) of UPHCs and 1/3 (33%) of public childbirth facilities.

Information on drugs was available from only two private clinics and three private childbirth facilities.

	Public facilities (n=23)		Private fac	cilities (n=23)
	UPHCs (n=20)	Childbirth facility (n=3)	Clinics (n=20)	Childbirth facility (n=3)
Functional registration counter	18 (90%)	0 (-)	11 (55%)	3 (100%)
Equipment				
✓ ECG machine	2 (10%)	1 (33%)	1 (5%)	3 (100%)
✓ Snellen's chart	6 (30%)	1 (33%)	1 (5%)	1 (33%)
✓ Ophthalmoscope	1 (5%)	1 (33%)	1 (5%)	1 (33%)
✓ Cardiopulmonary monitor	1 (5%)	2 (67%)	1 (5%)	1 (33%)

Table 5: Infrastructure and equipment available in health facilities

Lab equipment				
✓ Differential blood cell count machine	4 (20%)	2 (67%)	0 (-)	3 (100%)
✓ Colorimeter	11 (55%)	2 (67%)	0 (-)	2 (67%)
✓ Biochemistry analyzer	13 (65%)	1 (33%)	0 (-)	3 (100%)

There was a waiting area for patients available in all UPHCs, public and private childbirth facilities and in 16/20 (80%) of private clinics. Only public and private childbirth facilities had a functional labor room. All the UPHCs, public and private childbirth facilities and 10/20 (50%) of private clinics have easily accessible pathways. Toilets were available in all UPHCs, public and private childbirth facilities and in only 3/20 (15%) of private clinics.

Other infrastructure facilities such as a room for practicing yoga or meditation were available in 11/23 (48%) of public facilities but not available in private facilities. A room was available for nutrition demonstration in only [4/23 (17%) and 1/23 (4%)] and for counseling patients [3/23 (13%) and 1/23 (4%)] of public and private facilities, respectively.

Only 2/23 (9%) of public facilities had a park within 50 meters. None of the health facilities had a gym close by. Most [16/23 (70%) each], of both public and private facilities had an average of three pharmacies on the same street; [13/23 (56%) and 16/23 (70%)] had an average of 2 clinics on same street; [10/23 (43%) and 11/23 (48%)] of public and private facilities respectively had an average of one lab in the same street.

Limited number of public facilities (<25%) had provision of a UID for registration of patients, ordering of tests, prescriptions, referral, and follow-up (Table 6) and this was non-existent in almost all private facilities.

	YES	NO
UID for each patient		
✓ Registration	5 (22%) *	18 (78%)
 ✓ Ordering tests 	5 (22%) *	18 (78%)
✓ Pharmacy	4 (17%) **	19 (83%)
✓ Referring	5 (22%) *	18 (78%)
✓ Follow-up	5 (22%) *	18 (78%)
Reports sent to the		
government.	21 (91%)	2 (9%)
✓ Births	21 (91%)	2 (9%)
✓ Deaths	22 (96%)	1 (4%)
✓ Communicable diseases	, , ,	、 <i>、 、</i>

Table 6: Availability of health	information technology	at public facilities (n=23)
---------------------------------	------------------------	-----------------------------

Equipment for data entry		
✓ Computer	23 (100%)	-
 ✓ Online support 	23 (100%)	-
Passive data entry		
✓ Aggregate numbers	12(52%)	11 (48%)
✓ Facility data	23(100%)	-
 ✓ Community data 	23 (100%)	-
Registration with NDHM	2 (9%)	21 (91%)

**Kumbarakoppal, Bannimantap, Vishweswaranagar, Chamundipura, Giribhovipalya* ** *except for Kumbarakopall others are included*

All the public facilities (100%) had outsourced waste management to an external agency, while only 5/23 (22%) of private facilities reported outsourced waste management. Basic amenities for infection control such as wash basins, color coded bins in relevant places - lab and treatment area were available in all the public facilities, but this was not available in most of the private clinics.

Only 5/20 (25%) and 2/3 (67%) of UPHCs and public childbirth facilities had provided all staff welfare activities for their staff (Table 7).

Table 7: Staff welfare activities in public facilities (n=23)

	UPHCs (<i>n</i> =20)	Childbirth facility (n=3)
All below staff welfare services provided: 7 (30%)	5 (25%)	2 (67%)
 Immunisations offered – all four* ✓ Tetanus toxoid ✓ Typhoid ✓ Hepatitis B ✓ Covid 19 	6 (30%) 13 (65%) 9 (45%) 15 (75%) 17 (85%)	2 (67%) 3 (100%) 2 (67%) 3 (100%) 3 (100%)
Post exposure prophylaxis (PEP) available – 97%	12 (60%)	3 (100%)
Annual health check for staff available - 65%	19 (95%)	3 (100%)

Private childbirth facility- 2 (67%) offered all immunisations; offered annual health checks; offered PEP. Except for one private clinic, all others had only a doctor and sometimes a class D worker for cleaning the facility. Hence these details were not ascertained in the clinics.

C.1.2. Services provided by health facilities.

Table 8: Maternity and	l child health servic	es available at facilities
------------------------	-----------------------	----------------------------

	Publ	lic (n=23)	Priv	ate (n=23)
Clinical services	UPHCs (n=20)	Childbirth facility (n=3)	Clinics (n=20)	Childbirth facility (n=3)
 Maternity and neonatal services ✓ Antenatal care ✓ Postnatal care ✓ Labor ✓ Neonatal 	20 (100%) 20 (100%) 0 (-) 11 (55%)	3 (100%) 3 (100%) 3 (100%) 3 (100%)	7 (35%) 2 (10%) 0 (-) 2 (10%)	3 (100%) 3 (100%) 3 (100%) 3 (100%)
 Child health services ✓ Immunization ✓ Treatment- minor illnesses/problems ✓ Growth monitoring ✓ First aid for injuries ✓ Referral ✓ Emergency services 	20 (100%) 20 (100%) 19 (95%) 20 (100%) 19 (95%) 19 (95%)	3 (100%) 3 (100%) 3 (100%) 3 (100%) 3 (100%) 3 (100%)	3 (15%) 9 (45%) 3 (15%) 8 (40%) 2 (10%) 2 (10%)	2 (67%) 3 (100%) 3 (100%) 3 (100%) 2 (67%) 3 (100%)

Only 11/20 (55%) of UPHCs and 2/20 (10%) of clinics offered neonatal services. Private clinics provided limited maternity services 7/20 (35%) only offered antenatal care and 2 /20 (10%) offered postnatal care as seen in Table 8. Majority of the public facilities (>95%) offered all the child health services, while only a limited number of private clinics offered child health services such as immunisation 3/20 (15%); growth monitoring 3/20 (15%)! referral and emergency services 2/20 (10%).

The commonest reasons for referrals of children as cited by public facilities included:

- ✓ Severe injuries -9/23 (39%);
- ✓ Gastro-enteritis: diarrhoea, vomiting 6/23 (26%);
- ✓ Malnutrition: severe / moderate 5/23 (22%);
- ✓ Respiratory problems -4/23 (17%);
- ✓ Very high fever -3/23 (13%)
- ✓ Convulsions -3/23 (13%)

 \checkmark Skin problems – 3/23 (13%)

The commonest child health emergencies as elicited from public facilities included:

- ✓ Severe injuries -15/23 (65%);
- ✓ Gastro-enteritis -5/23 (22%);
- ✓ High fever -5/23 (22%);
- ✓ Respiratory problems -4/23 (17%);
- ✓ Malnutrition: severe / moderate 2 (9%)

Table 9: Reported adult health services of public and private facilities

	<i>Public</i> (<i>n</i> =23)		Private (n	(<i>n</i> =23)	
Adult clinical services	UPHCs	Childbirth	Clinics	Childbirth	
	(n=20)	facility (n=3)	(n=20)	facility (n=3)	
 ✓ Treatment of minor illnesses ✓ Provision of medical certificate ✓ First aid for minor injuries ✓ Minor surgical interventions (abscess/wound) 	20 (100%) 17 (85%) 20 (100%) 18 (90%)	3 (100%) 3 (100%) 3 (100%) 3 (100%) 3 (100%)	17 (85%) 2 (10%) 17 (85%) 12 (60%)	3 (100%) 3 (100%) 3 (100%) 3 (100%) 3 (100%)	
 ✓ Treatment of NCDs Diabetes Hypertension Cancer COPD 	20 (100%)	3 (100%)	11 (55%)	2 (67%)	
	20 (100%)	3 (100%)	11 (55%)	2 (67%)	
	4 (20%)	2 (67%)	1 (5%)	2 (67%)	
	16 (80%)	2 (67%)	6 (30%)	2 (67%)	
Mental healthThyroid problem	3 (13%) 0		1 (4%) 1 (4%)		
 ✓ Treatment of communicable diseases Tuberculosis Malaria Dengue Chikungunya Typhoid Leprosy STI All viral diseases 	20 (100%)	3 (100%)	9 (39%)	1 (33%)	
	14 (70%)	2 (67%)	6 (30%)	2 (67%)	
	4 (20%)	2 (67%)	6 (30%)	2 (67%)	
	4 (20%)	1 (33%)	0 (-)	0 (-)	
	5 (22%)	0 (-)	7 (35%)	1 (33%)	
	7 (35%)	1 (33%)	0 (-)	0 (-)	
	2 (9%)	0 (-)	0 (-)	0 (-)	
	0 (-)	0 (-)	3 (15%)	0 (-)	

Less than 35% of UPHCs were providing treatment of cancer, mental health, and thyroid problems as well as for dengue, chikungunya, typhoid, leprosy and STIs among communicable diseases while rest of adult health services were being provided by more than 70% (Table 9). Among the private clinics only 55% offered treatment for diabetes and hypertension and <40% offered treatment for COPD, and other communicable diseases such as tuberculosis, malaria, dengue, and typhoid.

The commonest emergency reported was injuries by both public and private [10/23 (44%) and 4/23 (17%)] facilities respectively (Table 10). Injury was also reported as the commonest reason for a referral by 14/23 (61%) of public facilities and 3/23 (13%) of private facilities.

Reported adult emergency and referral services	Public (n=23)	Private (n=23)
 ✓ Emergency services provided Injuries (burns/wounds/fractures/abscess) Road traffic accident cases Cardiac (heart attack/Low BP/high BP) Diabetes related (DKA/hypoglycemia) Respiratory (COPD/Asthma/Breathing difficulty) Bites (dog/snake) CNS related (epilepsy/unconscious) Suicide 	10 (44%) 9 (39%) 8 (35)% 5 (22%) 5 (22%) 3 (13%) 2 (9%) 1 (4%)	4 (17%) 3 (13%) 1 (4%) 1 (4%) 1 (4%) Not reported Not reported Not reported
 ✓ Referral in the last month? Injuries Cardiac cases Accident Bites CNS related Others (MLC/suicide) Respiratory related Renal cases Cancer cases 	14 (61%) 6 (26%) 5 (22%) 4 (17%) 3 (13%) 3 (13%) 1 (4%) Not reported Not reported	3 (13%) 3 (13%) 3 (13%) Not reported Not reported Not reported 3 (13%) 2 (9%) 1 (4%)

Table 10: Adult emergency and referral services reported by public and private facilities

Table 11 and Table 12 provides information on load of services and lab investigations performed for a month in public and private facilities. Table 11 provides information on reach of services by field health care workers such as the ANMs and ASHAs.

Table 11: Monthly load of services of public facilities based on record review of the last month

	Median per public facility for one monthUPHCs (n=20)Childbirth facility (n=3)		Median per private facili for one month*	
			Clinics (n=20)	Childbirth facility (n=3)
OPD registration-Morning	1447	1952		1140
ANC registration	26	353		17
Newborn registration as per ASHA	6	Not elicited	Ν	Not elicited
Postnatal registration as per ASHA	6	Not elicited	0	Not elicited
Outreach sessions	2	0	Т	0
Emergencies managed	3	7		Not validated

Acute malnutrition – referred to NRC	2	0	А	0
 Children treated ✓ Anemia ✓ Diarrhea 	$\begin{array}{c} 0 \\ 4 \end{array}$	0 1	V A I L	0 4
 Number of adults with ✓ TB on treatment ✓ TB – completed treatment ✓ Diabetes – New ✓ Diabetes – Old ✓ Hypertension – New ✓ Hypertension - Old 	13 1 5 94 4 134	21 2 1 73 2 68	A B L E	21 2 10 150 10 150
Referrals made ✓ Maternal ✓ Neonatal ✓ DM ✓ HTN ✓ Cancer ✓ Palliative	0 0 0 0 0 0	9 2 0 0 0 0 0		0 1 0 0 0 0

*OPD services/Services as prescribed under RCH-II, NH programmes, referral services, basic lab services, outreach services * only one or two facilities reported details; Record review performed either in May, June, July, August 2022.*

Table 12: Lab investigations performed in a month based on record review of the last month					
Lab investigations		per month in ic facilities	Median per month in private facilities		
	UPHCs (n=20)	Childbirth facility (n=3)	Clinics (n=20)	Childbirth facility (n=3)	
Pregnancy tests	10	40		10	
Hemoglobin	71	362		81	
Blood group and typing	19	145		15	
HIV	21	222	N O T A P P	28	
HBsAg	14	130		25	
VDRL	14	128		15	
Malaria smear test	384	176		8	
AFB sputum test	8	18	L	3	
Routine urine test	38	322	I C A B	60	
Blood sugar – RBS	169	709		51	
HbA1C*	5	NA	L E	10	
Lipid profile	0	0	Ľ	19	

Table 12: Lab investigations performed in a month based on record review of the last month

Thyroid test	0	0	38
Serum potassium and creatinine	0	0	25
Dengue	0	0	10

*Only 12/23 public facilities performed HbA1C. Record review performed either in May, June, July, August 2022.

Majority (87%) of the public facilities reported outreach services (Figure 5). None of these outreach services were reported by private facilities.




 Table 13: Median number of persons reached for health services at the community by

 UPHCs (n=20)

Presently registered	Currently registered Median** by ANMs	Currently registered Median** by ASHAs
Pregnant women	40	7
Newborns	-	6
Postnatal mothers	-	6
Adults with diabetes	171	211

Adults with hypertension	180	212
Adults on TB treatment	4	
Others (cancer/mental health)	5	2

*1 UPHC did not have an ASHA who could do the survey; **Surveyed in Apr or May 2022

Health workers – numbers against standards, roles, challenges in providing services

Table 14: Health workers filled against required Standards for UPHCs (n=20)

Health workers** [1000000 population approximately]	Required based on Standards	Filled	Gap (Filled posts from Standards)
Doctors (1 fulltime + 2 parttime)	20+60	21	0
Nurses (3/UPHC)	60	44	16 (27%)
ANMs (10000 population/ANM)	100	83	17 (17%)
ASHAs (2500 population/ASHA)	400	81	319 (80%)
Lab technician (1 per UPHC) *	20	22	0
Pharmacist (1 per UPHC) **	20	15	5 (25%)
DEO / M & E Unit (1 per UPHC)	20	8	12 (60%)
 Others (LDC / SDA) PH manager one per UPHC LHV 1 per UPHC 2 support staff per UPHC 	80 20 20 40	14	27 (34%)
Health Inspection Officer		39	
Class D worker (3 per UPHC)	60	39	21 (35%)
Counsellors			

*1 UPHC with 2 and 1 without a lab technician; ** 5 UPHCS with no pharmacist, acting pharmacists: 3 nurses and 2 medical officer

Table 14 shows the gap in HCPs both facility level and field based at the UPHCs. There was 80% gap of ASHAs based on standards required for the post.

Of the public facilities: two had gynaecologist and pediatrician on call; while of private facilities: three had gynaecologist, four had pediatricians, two had pediatric surgery doctors on call.

More than 90% of nurses and lab technicians were appointed on contract basis while 76% of doctors had permanent appointment (Table 15).

	Doctors* (n=21)	Nurses (n=46)	Lab technician (n=22)	Pharmacist (n=17)	ANM (n=84)	ASHA (n=81)
 Appointment ✓ On contract ✓ Through NGO ✓ Permanent 	3 (14%) 2 (10%) 16 (76%)	43 (93%) 0 (-) 3 (7%)	21 (95%) 0 (-) 1 (5%)	10 (59%) 0 (-) 7 (41%)	60(71%) 6 (7%) 18 (22%)	81 (100%) 0 (-) 0 (-)
Age in years ✓ (Mean±SD) ✓ Range	45.8±11.6 25-59	37.1±9.2 22-58	32.9±7.3 24-50	35.3±9.1 22-54	34.2±7.2 24-57	36.4±6.3 24-54
Sex ✓ Male ✓ Female *1 facility had 2 de	8 (38%) 13 (62%)	0 (-) 46 (100%)	9(41%) 13 (59)%	2 (12%) 15 (88%)	0 (-) 84 (100%)	0 (-) 81 (100%)

Table 15: Demographic variables of health workers from UPHCs (n=20)

More than 80% doctors and nurses had received supervision in the last 3 months, while 60% of them were monitored monthly by a senior.

Amongst doctors and nurses only 11 (48%) and 10 (30%) received training on Rashtriya Bal Swasthya Karyakram (RBSK); 12 (52%) and 8 (35%) received training on Rashtriya Kishor Swasthya Karyakram (RKSK) respectively (Table 15). These programs are focused on the overall quality of life and health of children and adolescents.

One of each health workers was selected for assessing training received (Table 16-Table 18), their roles (Table 19-20) and challenges faced in providing required services (figure 15-16) are presented further.

Table 16: Training received as reported by doctors and nurses from public facilities

Areas of training received in the last 5 years	Doctors (n=23)	Nurses (n=23)
SBA	8 (35%)	12 (52%) *
New-born	16 (70%)	23 (100%) **
RBSK	11 (48%)	10 (30%)
Infection control	15 (65%)	21 (91%)
Family planning	17 (74%)	21 (91%) *

 Non-Communicable Diseases (NCDs) – YES ✓ Diabetes ✓ Cancers ✓ Hypertension (HTN) ✓ Mental Health (MH) problems ✓ Injuries ✓ COPD 	17 (74%) 17 (74%) 17 (74%) 17 (74%) 16 (70%) 14 (61%) 13 (57%)	23 (100%) 23 (100%) 23 (100%) 23 (100%) 19 (83%) 18 (78%) 0 ()
RKSK	12 (52%)	8 (35%) **
Communicable Diseases (CDs)- ✓ Tuberculosis (TB) ✓ Dengue ✓ HIV ✓ Diarrheal diseases	18 (78%) 17 (74%) 15 (65%) 15 (65%)	16 (70%) 12 (52%) 0 (-) 11 (48%)

*3 from public childbirth facility received training; ** 1 from public childbirth facility

Table 17: Training reported to have been received by lab technicians and pharmacies at all public facilities (n=23)

Areas of training received in the last 5 years	Lab technicians (n=23)
Malaria / National Vector Borne Disease Control Program (NVBDCP)	18 (78%)
Tuberculosis	13 (57%)
NUHM	8 (35%)
Biomedical waste management	4 (17%)
Non-Communicable Diseases	4 (17%)
IHIP (Integrated health information platform)	3 (13%)
No training	1 (4%)
Areas of training received in the last 5 years	Pharmacists (n=23*)
Electronic Vaccine Intelligence Network (eVIN) app/Karnataka State Medical Supplies Company (KSMSC)	11 (47%)
Vaccine storage	5 (22%)
Nikshay (Personal Protective Equipment and products)	3 (13%)
Biomedical waste management	1 (4%)
Integrated Disease Surveillance Project (IDSP)	1 (4%)
Did not respond to question	2 (9%)

*5 UPHCs and 1 public childbirth facility did not have qualified pharmacists. The nurse / doctor acting as pharmacists gave the interview.

Most of the lab technicians 18 (78%) had training on NVBDCP and on tuberculosis 13 (57%). While amongst pharmacists 11 (55%) mentioned they received training on eVIN (Table 17).

Two (9%) of the lab technicians mentioned that they had stock-out of lab supplies in the last three months and their equipment were not functional. While among the pharmacists and acting pharmacists (4 nurses and 2 medical officers), 10/23 (44%) mentioned that they had medication stock outs. These included drugs for treatment of diabetes and hypertension.

Area of training in the last 5 years	ANMs* (n=21)	ASHA** (n=20)
SBA	14 (67%)	Module1-20 (100%)
Newborn	19 (90%)	Module 2- 20 (100%)
RBSK	8 (35%)	Module 3 – 20 (100%)
Infection control	12 (57%)	Module 4 – 18 (90%)
NCDs ✓ Diabetes ✓ Cancers ✓ Hypertension ✓ Mental health ✓ Injuries	16 (76%) 17 (81%) 16 (76%) 13 (62%) 16 (76%)	Module 5 – 13 (65%) Module 6 – 10 (50%) Module 7 – 4 (20%) All Modules – 1 (5%)
RKSK	7 (33%)	
Communicable diseases ✓ TB ✓ Dengue ✓ Diarrheal diseases	17 (81%) 17 (81%) 17 (81%)	*No ANMs and ASHAs in 2 public childbirth facilities **No ASHA in one UPHC

Table 18: Training reported to have been received by ANMs and ASHAs in public facilities

Most of the ANMs (Table 18) reported to have been trained on newborn care [19 (90%)], on cancers [17 (81%)], on Diabetes and hypertension [16 (76%)] and on communicable diseases – TB, dengue, diarrheal diseases [17 (81%)].

Of the ASHAs (Table 18), most (>90%) had training on Module 1 that introduced them to their roles and responsibilities, Module 2 on MCH, Module 3 on family planning, contraceptives, Reproductive and Sexual Health (RSH) including Adolescent RSH, Module 4 on National health programs including AYUSH and management of minor ailments. While training on Module 5 which was on knowing self, human rights, leadership, skills-communication, decision-making, negotiation, coordination; Module 6 on MNH skills to save lives; Module 7 Neonatal and Child health skills to save lives was received by 65%, 50% and 20% respectively.

Doctors' role at public facilities	Frequency of performance							
	Most often		Less often		Rarely		Blank /Not responded	
	а	b	а	b	а	b	а	B
Clinical management of patient	19	1	0	0	0	0	1	2
Visit subcenter	12	0	5	0	2	0	1	3
Preparation of operational plans	12	0	2	0	4	0	2	3
School visits	10	0	4	0	4	0	2	3
Supervision of nurses and health workers	18	1	0	0	1	0	1	2
Training of staff	16	1	1	0	2	0	1	2
Intersectoral coordination	12	1	5	0	2	0	1	2
Supervision of administrative works	16	1	2	0	1	0	1	2
Coordination with AYUSH*	8	0	1	0	2	0	9	3
Field visits to understand local problems	15	0	2	0	2	0	1	3

Table 19: Role of doctors at public facilities (n=23)

a=UPHCs (n=20); b= public childbirth facilities (n=3); *AYUSH: Ayurveda, Siddha, Homeopathy medicine

The most often roles of doctors as seen in Table 19 at public facilities included clinical management of patients 19/23 (82%), supervision of staff and other health workers 18/23 (78%), supervision of administrative work and training of staff – 16/23 (70%). All 23 of private facility doctors only mentioned clinical management as their main role.

Staff nurses' roles at public facilities	Frequency of performance based on rank order (n=23)							k order	
	Most often					ely	Not responded / Blank		
	a	b	a	В	a	b	а	b	
Clinical management of patient	20	3	0	0	0	0	0	0	
Conduct of MCH clinics	17	1	2	0	0	3	0	0	
Outreach camps /School health services*	9	0	5	0	4	1	2	2	
Equipment – functional and maintained	17	3	3	0	0	0	0	0	
Supervision of other HCPs*	5	0	0	0	5	0	10	3	
Conduct deliveries	0	3	0	0	0	0	20	0	

Table 20: Role of nurses at public facilities (n=23)

Cleanliness and IC in facility	14	3	4	0	0	0	1	0
Educational activities – Facility / Community*	17	3	3	0	0	0	0	0
Attending meetings with others – ANMs / ASHAs*	9	0	1	0	5	0	5	3

• Not relevant for private health facilities and only two facilities had a nurse.

• *a*=*UPHCs*; *b*=*public childbirth facility*

The most often roles of nurses at public facilities included clinical management 20/23 (87%); Educational activities, conduct of MCH clinics and maintenance of equipment 17/23 (73%). Only 2 private health facilities had nurses and they reported clinical management, conduct of MCH clinics, management of women during childbirth and supervision of the cleanliness of the facility were their main roles.

More than 91% of ANMs and ASHAs reported to have been supervised and 100% were monitored by a senior monthly.

C.2. Profile the community morbidity status, healthcare seeking, and costs incurred for selected acute and chronic conditions in urban wards of Mysuru city

Survey was carried out in 6007 households of Mysuru urban, comprising of 21576 individuals, having 51.2% females. Table 1 represents the summary of the demographic characteristics of the surveyed population. Among 6474 women in the reproductive age group, 100 (1.6%) mothers were currently pregnant and pregnancy in the last 3 years were 563 (8.6%). Among 11978 individuals aged above 30 years, 17.3% of were either diabetes or hypertension.

C.2.1. Demographic profile of the community

Parameters	N (%)
Number of HH	6007
Individuals	21576
Current pregnancy (18-49)	100 / 6474 (1.6%)
Pregnancy in the last 3 years $(18 - 52 \text{ years})$	563 / 6555 (8.6%)
Reproductive age group women $(18 - 49 \text{ years})$	6474 (30.0%)
Children below 5 years	1262 / 21576 (5.8%)
Children sick below 5 years	256 / 1262 (20.3%)
Either Diabetes or hypertensives (age > 30 years)	2078 / 11978 (17.3 %)

 Table 21: Demographic characteristics of the study population

Out of total population surveyed, 89.2% belonged to less than 60 years of age and 55.5% of them were above 30 years of age. The age distribution by gender of the study population is given in Table 22. There were 5.8% (n=1262) of children below 5 years of age, among them 25% were sick in the past 1 month (Table 23).

Age group (in	Male, $N = 10528$	Female, $N = 11048$	Total, N (%)
years)	(48.8%)	(51.2%)	
0-5	680 (6.5%)	582 (5.2%)	1262 (5.8%)
6-20	2178 (20.7%)	2109 (18.9%)	4287 (19.8%)
21-35	2757 (26.2%)	3199 (28.7%)	5876 (27.5%)
36-50	2593 (24.6%)	2764 (24.8%)	5358 (24.7%)
51-65	1574 (14.9%)	1747 (15.7%)	3321 (15.3%)
≥66 years	746 (7.9%)	727 (6.5%)	1472 (6.8%)

Table 22: Descriptive statistics of age group by gender of the population

Age	Male, N = 680 (%)	Female, N = 582 (%)	Total, N = 1262 (%)
Below 1 year	55 (61.1%)	35 (38.9%)	90 (7.1%)
1 – 5 years	625 (53.3%)	547 (46.7%)	1172 (92.8%)
Total	680 (53.9%)	582 (46.1%)	1262

Socio-demographic characteristics such as marital status, education and occupation of the population is presented in Table 24. Among 16907 individuals \geq 18 years of age, 69.2% were currently married. Among 20298 individuals \geq 6 years of age, 22% had completed graduation and above and one fourth of them had completed high school. Majority (~60%) of the females were home maker. More than one fifth of the population (21.6%) were employed either in government or private sector, approximately 6% of the individual were unemployed during survey period.

Table 24: Demographic	parameters of the surveyed	population
Table 24. Demographic	parameters of the surveyed	population

Demographic Parameters	Male, N (%)	Female, N (%)	Total, N (%)
Marital Status (> 18 years)	8084 (76.8%)	8823 (79.9%)	16907 (78.4%)
Never married	1722 (21.3%)	1082 (12.3%)	2804 (16.6%)
Currently married	5752 (71.2%)	5948 (67.4%)	11700 (69.2%)
Widowed/Separated/Divorced	290 (3.6%)	1589 (18.0%)	1879 (11.1%)
Education (≥6 years)	9843 (93.5%)	10455 (94.6%)	20298 (94.1%)
Illiterate	866 (8.8%)	1589 (15.2%)	2454 (12.1%)
Primary and middle	2300 (23.3%)	2480 (23.7%)	4780 (23.5%)
High school	2666 (27.1%)	2793 (26.7%)	5459 (26.9%)
PUC/ Class 11&12	1556 (15.8%)	1528 (14.6%)	3084 (15.2%)
Graduate	2075(21.1%)	1713 (16.4%)	3788 (18.7%)
Postgraduate and above	380 (3.9%)	353 (3.4%)	733 (3.6%)
Occupation	10472	10975	21447
Govt	358 (3.4%)	115 (1.0%)	473 (2.2%)

Private	2594 (24.8%)	807 (7.4%)	3401 (15.9%)
Business and Self employed	1741 (16.6%)	206 (1.8%)	1947 (9.1%)
Daily wages	1703 (16.3%)	396 (3.6%)	2099 (9.8%)
Retired	541 (5.2%)	94 (0.9%)	635 (3%)
Student	2278 (21.8%)	2224 (20.3%)	4502 (21%)
Home maker	485 (4.6%)	6466 (58.9%)	6951 (32.4%)
Unemployed	772 (7.4%)	667 (6.1%)	1441 (6.7%)

Table 25: Household Characteristics of the surveyed population

Household characteristics	N (%)
Religion	6007
Hindu	4656 (77.5%)
Muslim	1227 (20.4%)
Christian	110 (1.8%)
Others	14 (0.2%)
Caste	6007
SC/ST	851 (14.4%)
OBC	2868 (48.5%)
General	2288 (38%)
BPL card shown	4170 (69.4%)
House ownership - Owned	4013 (66.8%)
House Type – Pucca/Semi pucca/Kutcha	4904 (81.6%) / 1024 (17%) / 79
	(1.3%)
Drinking water -Within the premises/ Near the	5015 (83.5%) / 565 (9.4%) / 420
premises / Away	(7.0%)
Toilet – Improved not shared / Shared facility /	3538 (58.9%) / 2419 (40.3%) / 50
Unimproved & no facility	(0.8%)
Solid waste segregated at source- Yes	5671 (94.4%)
Cooking fuel - LPG/electricity	5955 (99.1%)
Fridge availability	3955 (65.8%)
Computer (laptop) with/ without internet / not	441 (7.3%)/158 (2.6%) / 5408 (90%)
available	
Motor vehicle 4 -wheeler	518 (8.6%)
3-wheeler	237 (3.9%)
2-wheeler	5178 (86.2%)
Phone Ownership Smart / Land phones	5584 (93%) / 1971 (32.8%)

Table 25 showed the household characteristics of the surveyed population. Most of the surveyed population were Hindu by religion (77.5%) and OBC by caste (48.5%). BPL card was shown by 70% of the surveyed households. More than 2/3rd of the households was having their own houses. About 82% of the surveyed house were of Pucca in nature. Most of them had drinking water facility within their premises (83.5%) and improved toilet facility (99.0%). Majority of the segregated solid waste at source (94.4%) and used LPG as cooking fuel (99.1%) and these data were comparable to NFHS-5. Refrigerator was available in more than 2/3rd of the households. About 90% of the

households were not having either computer or laptop. Most of the households had two wheelers (86.2%) and average of two smart phones per household.

C.2.2. Health Insurance (Table.26):

Majority of the households' members did not have any kind of health insurance (70.5%) which was like what was observed in NFHS -5 data (73.8%). Among those who were having health insurance Ayushman Bharat/ Arogya Karnataka was the most reported followed by privately purchased health insurance.

Health Insurance	Yes, N (%)	
Type of Health Insurance		
Ayushman Bharat/Arogya Karnataka	989 (16.5%)	
Employees state Insurance scheme / CGHS	466 (7.8%)	
Other privately purchased health insurance	446 (7.4%)	
Medical reimbursement from employer	34 (0.6%)	
No insurance	4072 (67.8%)	
Insurance used in last 1 year	26	

Table 26: Health in	nsurance statistics	of the s	urveved i	opulation
I WOIC BOT HICKICH H			ar , c, ca	Jopanation

Health Insurance	Male, N (%)	Female, N (%)	Total, N (%)	
Type of Health Insurance				
Ayushman Bharat/Arogya Karnataka	1523 (14.5%)	1558 (14.1%)	3081 (14.3%)	
Employees state Insurance scheme / CGHS	783 (7.4%)	744 (6.7%)	1527 (7.1%)	
Other privately purchased health insurance	788 (7.5%)	792 (7.2%)	1580 (7.3%)	
Medical reimbursement from employer	59 (0.6%)	52 (0.5%)	111 (0.5%)	
No insurance	7325 (70.1%)	7902 (71.6%)	15227 (70.5%)	
Insurance used in last 1 year	26	28	54 (0.2%)	

C.2.3. Lifestyle characteristics (Table 27):

Tobacco consumption in both smoking and chewing form was observed in 3% of the population respectively among individuals aged above 18 years. Alcohol consumption was reported in 4.4%

of the individual's majority being males. Majority of the survey individuals consumed fruits (61.5%) and vegetables (92.8%) for more than 4 days a week. Nealy half of the surveyed individuals reported that to use salt at the time of food preparation. More than ¹/₄ the of the household members were doing brisk walk for more than 4 days a week. Considering eating fruits, vegetables and walking for more than 4 days a week and none of the habits of using tobacco and alcohol as healthy habits, only 13% of the surveyed individuals were noted to be having healthy lifestyle. Only 5% of the individual noted to be part of some voluntary organization.

Age group (>18yrs)	Male	Female	Total
	N = 7929 (%)	N = 8662 (%)	N = 16591 (%)
Smoking tobacco	470 (5.9%)	69 (0.8%)	539 (3.2%)
Chewing tobacco	387 (5.5%)	74 (0.9%)	461 (3.1%)
Alcohol	608 (7.7%)	117 (1.4%)	725 (4.4%)
Fruit Eating>=5	4346 (61.5%)	4846 (61.5%)	9192 (61.5%)
days/week			
Vegetable eating>=5 days/week	6584 (93.2%)	7304 (92.6%)	13888 (92.8%)
How often Add salt			
Always / Often	51 (0.8%)	45 (0.5%)	96 (0.6%)
Sometimes / Rarely	869 (12.3%)	870 (11.0%)	1739 (11.7%)
Never	2624 (37.1%)	2988 (37.9%)	5612 (37.5%)
At food preparation	3522 (49.9%)	3980 (50.5%)	7502 (50.2%)
Brisk walk	1844 (26.1%)	1970 (25%)	3814 (25.5%)
Eye check up on own	1274 (18.1%)	1522 (19.3%)	2796 (18.7%)
Unhealthy lifestyle	1037 (13.1%)	161 (1.9%)	1198 (7.2%)
Healthy lifestyle	1096 (13.8%)	1159 (13.4%)	2255 (13.6%)
Member of voluntary	N = 7066	N = 7884	N = 14950
organization			
Mahila Arogya Samiti	24 (0.3%)	50 (0.6%)	74 (0.5%)
Self-Help Group	71 (1%)	186 (2.4%)	257 (1.7%)
Other (Dharmasthala/Ujjivana /Gramina kuta)	69 (1%)	298 (3.8%)	367 (2.5%)
None	6902 (97.7%)	7350 (93.2%)	14252 (95.3%)

Table 27 Lifestyle characteristics (in \geq 18 years age group) of the surveyed population

C.2.4. 2-weeks morbidity:

Around 7% (n-1490) of the surveyed population reported to have some illness in the past 2 weeks (Figure 6). Almost half of them sought treatment at health facility. Among those not seeking treatment at health facility, majority reported using home remedies or medicines available at home.




Figure 6: Types of ailments in the surveyed population in the last 2 weeks

Health seeking behaviour (Table 28.1)

Public health facility was utilized by 32.5% of the individual who sought treatment and remaining went to private health facility (67.5%). Allopathy system of medicine was the most utilized system of medicine. One third of the patients were advised investigations and 84% were prescribed medicines for their ailments.




Figure 7: Place of health care for different ailments in the surveyed population in the last 2 weeks

Mode of travel: Most frequently used mode of travel for health seeking to health care facility was two-wheeler (38.0%) followed by walking (23.7%). Median distance to reach health care facility was around 2 km for both public and private facility. The reported time to reach the health facility

for illness was less than 30 minutes for most of them. Similarly, time to consult the doctor was also reported to be less than 30 minutes for majority of the individuals. Almost $2/3^{rd}$ (60.4%) of the patients preferred morning hours as time of consultation.

Cost of care analysis for acute ailments (Table 28.2)

The median consultation cost reported was 0 (0, 100) and 200 (100, 300) in public and private health care facility respectively. Almost 40% of those consulting in public facility did not pay any consultation fees. The median cost of investigations was 65 (10, 520) and 100 (10, 800) for public and private health care facility respectively. Significantly higher median cost was spent on medicines in private facility [290 (150, 500)] as compared to public [110 (0, 425)]. Travel cost was similar for both health facility [Rs. 50 (10, 100) Vs 75 (50, 127)]. Majority of the individuals reported to use either income or saving to manage their routine medical expenses.

Table 28.1: Health	seeking behaviour	characteristics in	n 2	weeks	morbidity	(for th	ne first
episode)							

2-week Morbidity	N (%)
Illness in the last 2 weeks (n = 21576)	1490 (6.9%)
Sought treatment at health facility	740 (49.7%)
0-5	57 (7.7%)
6-20	108 (14.6%)
21-35	95 (12.8%)
36-50	206 (27.8%)
51-65	160 (21.6%)
≥66	114 (15.4%)
Treatment received from health facility	N = 740
Govt	241 (32.5%)
Private	499 (67.5%)
Medicine system	N = 740
Allopathy	710 (96.0%)
Ayurveda	12 (1.6%)
Homeopathy	13 (1.8%)
Others	5 (0.6%)
At least 1 ailment	1490 (6.9%)
2 or more ailment	34 (0.2%)
Investigations advised	247 (35.7%)
Medicines prescribed	603 (84%)
Mode of travel	N = 720
Walk	171 (23.7%)
Bus	68 (9.4%)
Auto and others	129 (17.9%)
2-wheeler	273 (38%)
Car	79 (11%)




Figure 8: Distribution of individuals who sought treatment at health facility by age group




Figure 9: Distribution of individuals according to type of facility from where they received treatment

	Govt	Pvt	Total
Distance to HF median (IQR) in	2 (1, 4)	2.5 (1, 5)	2 (1, 4)
Km			
Time to reach the facility			
Less than 30 minutes	406 (56.7%)	135 (56.7%)	271 (56.7%)
30 minutes - 1 hour	268 (37.4%)	85 (35.7%)	183 (38.3%)
> = 1 hour	41 (5.9%)	18 (7.3%)	24 (5%)
Time taken to consult the doctor			
Less than 30 minutes	361 (51.4%)	112 (46.9%)	249 (53.8%)
30 minutes - 1 hour	250 (35.6%)	74 (31%)	176 (38.0%)
> = 1 hour	91 (13%)	53 (22.2%)	38(8.2%)
Time of consultation			
Morning	451 (67.7%)	187 (81.7%)	264 (60.4%)
Afternoon	80 (12%)	29 (12.7%)	51 (11.7%)
Evening	117 (17.6%)	7 (3.1%)	110 (25.2%)
Night	18 (2.7%)	6 (2.6%)	12 (2.7%)

Table 28.2: 2 weeks morbidity – Distance, time and cost involved (for the first episode of ailment)

Ailment 1	Govt	Pvt	Total
Consultation cost in Rs	0 (0, 100)	200 (100, 300)	110 (10, 250)
Consultation amount not paid	39.8%	6.3% (Rs 10)	
Testing cost in Rs	65 (10, 520)	100 (10, 800)	100 (10, 520)
Drugs cost in Rs	110 (0, 425)	290 (150, 500)	200 (100, 500)
Cost spent on travel in Rs	50 (10, 100)	75 (50, 127)	60 (30, 100)
Managing routine medical	Govt	Pvt	Total
expenses	N = 241	N = 477	N = 718
Income	97 (40.2%)	186 (38.9%)	283 (39.4%)
Savings	151 (62.7%)	306 (64.2%)	477 (66.4%)
Borrowed	1 (0.4%)	9 (2.0%)	10 (1.4%)
Selling property / jewelry	3 (1.2%)	2 (0.4%)	5 (0.7%)
Insurance	3 (1.2%)	2 (0.4%)	5 (0.7%)

C.2.5. Maternal health:

Among 6474 women in the reproductive age group, 100 (1.6%) mothers were currently pregnant and pregnancy in the last 3 years were 563 (8.6%).

Antenatal care:

About 46% and 44% of the mothers preferred public and private health facility for their antenatal care respectively. More than 2/3rd of the ANC mothers showed their thai card during interview and very few (2.0%) were reported to have missed one of the ANC visit to health facility. Two-wheeler (40.0%) was the preferred most of transport for the ANC care. The median cost spent on ANC care was 2900 (1125, 5000) for mother utilizing public health facility. In contrast, higher median cost of 15000 (5500, 20000) was the amount spent in private health facility. Almost of them reported to use either income or saving to manage their ANC care medical expenses. The median (9) satisfaction score was similar for both public and private.

Childbirth care:

Public health facility was utilized for Childbirth care by 55% of the women. The reported mode of Childbirth was LSCS and other form of assisted Childbirth in 47.6% of the mothers which was comparable to NFHS-5 data of Mysuru city. Cash transfer benefits like Janani Suraksha Yojana were availed by 45.3% of the delivered mothers. Car (64.5%) the most commonly mode of transport for Childbirth. Only 10 mothers reported to have some complications during Childbirth for which majority were referred and went to public health facility (60%). Significant difference was noted in the median cost spent for Childbirth between public [5000(4500, 10000)] and private health facility

[50000 (25000, 60000)]. Around 7% of them borrowed/sold property/ jewellery to manage their Childbirth care medical expenses. The median (9) satisfaction score was similar for both public and private.

Postnatal care:

Postnatal care data was available on 547 mothers. Slightly more than half of them (54.8%) availed PNC care in public health facility. Cash transfer benefits like Janani Suraksha Yojana were availed by 42.8% of the delivered mothers. Car (52.9%) the most commonly mode of transport for PNC services. Complications was reported in 6 mothers and majority (80%) were referred and went to private health facility. The median cost spent for PNC care in public [3000(2000, 5000)] and private health facility [20000 (8000, 50000)]. Almost of them reported to use either income or saving to manage their PNC care medical expenses. The median (9) satisfaction score was similar for both public and private.

Maternal Health	ANC	CHILDBIRTH	PNC
	N = 100 (%)	N = 563 (%)	N = 547 (%)
Place of care			
Govt	46 (46%)	304 (54.7%)	300 (54.8%)
Private	44 (44%)	250 (45%)	246 (45%)
Traditional	1 (1%)	2 (0.4%)	1 (0.2%)
Primary respondent was not aware	9 (9%)	7 (1.2%)	0 (0%)
Thai card showed	77 (77 %)	512 (92.8%)	489 (89.4%)
Type of Childbirth			
Normal		289 (52.5%)	
		Govt: 216	
		(74.7%)	
		Private: 73	
		(25.3%)	
LSCS & others		262 (47.6%)	
		Govt: 86 (33%)	
		Private: 175	
		(67%)	
JSY / other cash transfer		248 (45.3%)	227 (42.8%)
Mode of transport	ANC	CHILDBIRTH	PNC
Walk	10 (11.1%)	6 (1.1%)	28 (5.2%)
Bus	8 (8.9%)	30 (5.4%)	28 (5.2%)
Auto and others	12 (13.3%)	122 (22.1%)	113 (20.9%)
2-wheeler	36 (40%)	38 (6.9%)	86 (15.9%)
Car	24 (26.7%)	356 (64.5%)	286 (52.9%)
Missed ANC visits	2 (2%)		
Complications (mother)	0 (0%)	10 (1.8%)	6 (1.1%)
Type of hospital went	NA	10	5

 Table 29.1: Maternal health: Health seeking behaviour characteristics

Govt	6 (60%)	1 (20%)
Private	3 (30%)	4 (80%)
None	1 (10%)	0 (0%)

	ANC	CHILDBIRTH	PNC
HH spend [Median	5000 (2000, 15000)	10000 (5000, 50000)	5000 (2000, 20000)
(IQR)]			
Govt	2900 (1125, 5000)	5000 (4500, 10000)	3000 (2000, 5000)
Pvt	15000 (5500, 20000)	50000 (25000, 65000)	20000 (8000, 50000)
Routine medicine	N = 100	N = 563	N = 547
expenses			
Income	35 (35%)	224 (39.8%)	226 (41.3%)
Savings	66 (66%)	385 (68.4%)	373 (68.2%)
Borrow from family	2 (2%)	15 (2.7%)	12 (2.2%)
Selling property /	0	15 (2.7%)	9 (1.6%)
jewelry			
Insurance	1 (1%)	6 (1.1%)	3 (0.5%)
Satisfaction score			
Median	8 (8,9)	9 (8,9)	8.5 (8,9)

 Table 29.2: Maternal health: Cost of care and satisfaction score

C.2.6. Child Health:

The mean age of children (\leq 5 years) was 2.7 ± 1.5 years. One fourth of the children (25%) were reported to be sick in the last 1 month. The most common illness reported was Acute Diarrhoeal Disease (ADD) (67.6%) followed by Acute Respiratory Infection (ARI) (35.5%). For both the ailments, approximately equal proportion of households availed care from public and private health facilities. The preferred reasons for choosing public health facility were less / free of cost followed by trust in doctor and nearby distance which was similar for both ARI and ADD. Trust in doctor, timely service, and all facilities at one place were the reasons reported in favour of private health facility. Time spent for consultation was less than 30 minutes in public health facility, and 30 – 60 minutes in private health facility. Child hospitalization rate was 13.2% and 4.0% for ARI and ADD respectively. The median cost spent on treatment for ARI in public was 128 (87, 335) and in private health facility and 145(100, 200) in private health facility. The median (9) satisfaction score was similar for both public and private.

Child Health (age ≤ 5	N = 1065			
years)				
Mean age	2.7 ± 1.5 year	s		
Currently sick in the last	256 (24%)			
1 month				
Ailment				
ARI	91 (35.5%)			
ADD	173 (67.6%)			
Others (Speech and	2 (0.8%)			
hearing issues; cleft lip)			1	
		ARI	A	DD
Place of treatment				
Govt		(50.5%)	,	50.6%)
Private	45 ((49.5%)	85 (4	49.4%)
Child hospitalized	12	(13.2%)	7 (4	4.0%)
Primary reason	Govt	Pvt	Govt	Pvt
Close by	15 (23.4%)	1 (4%)	14 (8%)	6 (3.4%)
Less cost/free of cost	29 (45.3%)	2 (8%)	117 (66.9%)	8 (4.5%)
Trust / good doctor	16 (25%)	15 (60%)	39 (22.3%)	81 (45.8%)
Timely service	3 (4.7%)	5 (20%)	3 (1.7%)	56 (31.6%)
All facility at one place	1 (1.5%)	2 (8%)	2 (1.1%)	26 (14.7%)
Time spent on	Govt	Pvt	Govt	Pvt
consultation				
<30 minutes	14 (53.8%)	5 (29.4%)	57 (65.5%)	37 (43.5%)
30 minutes -1 hour	7 (26.9%)	9 (52.9%)	24 (27.6%)	38 (44.7%)
1-2 hours	5 (19.2%)	3 (16.8%)	6 (6.9%)	10 (11.8%)
Money spent on	128 (87,	500 (185,	60 (50, 100)	145 (100, 200)
treatment	335)	10500)		
Median satisfaction score	8 (8, 9)	9 (8, 9)	8 (8, 9)	9 (8, 9)

Table 30: Child health: Health seeking behaviour, and satisfaction score

Child immunization:

Table 31 shows the proportion of children (<=24 months) completely immunized was 83.0%. About 73.4% of them received immunization from public health facility. Primary reason for preferring public health facility was less/free of cost (81.9%) followed by trust in doctor (57.9%).

Child immunization Done	<= 24 months)	N = 463 / 553	(83.7%)
Place			
Govt		340 (73.4%)	
Private		123 (26.6%)	
Primary reason	Govt		Pvt
Close by	92 (27%)		12 (9.7%)

Table 31: Health care seeking characteristics for child immunization

Less cost/free of cost	303 (89.1%)	9 (7.4%)
Trust / good doctor	197 (57.9%)	103 (83.7%)
Timely service	27 (7.9%)	61 (49.6%)
All facility at one place	22 (6.5%)	32 (26%)
Time spent for immunization		
<30 minutes	159 (47%)	56 (45.5%)
30 minutes -1 hour	154 (45.6%)	58 (47.2%)
1 – 2 hours	25 (7.4%)	9 (7.3%)

Non-Communicable diseases (Diabetes Mellitus and Hypertension):

Data pertaining to diabetes and hypertension was assessed in individuals aged more than 30 years. The reported prevalence of either diagnosed diabetes or hypertension was 17.3%, (15.9% in males / 18.7% in females). Both diabetes and hypertension were presented in 7.5% of the individuals (6.5% males/ 8.7% females). Diabetes and Hypertension alone was reported in 12.8% and 15.0% respectively.

For both diabetes and hypertension, approximately 65% of the individuals preferred private health facility and the allopathy was the preferred system of medicine (99.0%). The reason for preferring private health facility were trust in doctor (80.0%) followed by timely service (50.0%) and all facility at one place (27.0%). Around 3 fourth of the patients are on regular treatment for their medical conditions. Higher proportion of both diabetes (70.3%) and hypertension (65.9%) patients preferred private facility for buying medicine routinely. The regular fasting blood sugar check among diabetic individuals was done in private health facility (69.5%).

About 7% of the diabetic individuals and 3.8% of the hypertensives were told to have some complications by the consultant. Most of them were referred to private health facility for their complications. Eye, kidney, cholesterol, and ECG check up in the last 1 year was reported to be done in 63.7%, 69.5%, 69.4% and 70.9% of the diabetic individuals respectively. Among hypertensives, Eye, kidney, cholesterol, and ECG check up in the last 1 year was reported to be done in 54.1%, 44.5%, 46,7% and 56.8% respectively. Table x provides the data on recent blood sugar levels and the recorded BP at the time of survey. Like other illnesses, income and savings were reported to be the commonest mode of managing routine medical expenses in NCD patients. The median satisfaction score was 8 (8,9) in both public and private health facility.

Characteristics	DM (N= 1313)		HTN (N=1539)	
Gender distribution	Male	Female	Male	Female
	619 (47.1%)	694 (52.9%)	655 (42.6%)	884 (57.4%)
Routine visit				
Govt	415 (31.6%)		532 (35.4%)	

Table 32.1: Non-Communicable Diseases: Health seeking behaviour characteristics

Private	898 (68.4%)	969 (64.6%)
System of medicine		
Allopathy	1291 (98.5%)	1475 (98.7%)
Ayurveda	16 (1.2%)	12 (0.8%)
Others	4 (0.3%)	8 (0.6%)
Reasons for preference Govt		
Close by	80 (19.3%)	99 (18.6%)
Less cost/free of cost	423 (71.3%)	554 (71.8%)
Trust/Good doctor	246 (59.3%)	294 (55.3%)
Timely service	55 (13.2%)	62 (11.6%)
All facility at one place	51 (12.3%)	48 (9%)
Reasons for preference		
Private		
Close by	132 (14.9%)	123 (12.7%)
Less cost/free of cost	94 (6.3%)	103 (6%)
Trust/Good doctor	723 (80.5%)	789 (81.4%)
Timely service	420 (46.8%)	481 (49.7%)
All facility at one place	248 (27.6%)	257 (26.5%)
Last 6 months number of		
doctors consulted		
1 doctor	741	831
2 doctors	231	245
3 or more	58	49
Regular treatment	962 (73.4%)	1132 (76.1%)
Last 7 days missed taking		
medicine: For 1-2 days	117	160
3 and more days	26	20
Routine medicine	1301	1481
Govt	386 (29.7%)	505 (34.1%)
Private	915 (70.3%)	976 (65.9%)

Table 32.2: Non-Communicable Diseases: Details of investigations and complications

	DM (N = 1313)	HTN (N = 1539)
Eye checkup in last 1 year	827 (63.7%)	775 (54.1%)
Kidney checkup in last 1 year	902 (69.5%)	637 (44.5%)
Cholesterol checkup in last 1	902 (69.4%)	669 (46.7%)
year		
ECG checkup in last 1 year	920 (70.9%)	813 (56.8%)
Fasting blood sugar	N = 1259	
Govt	390 (31%)	
Private	869 (69%)	
Complications doctor told	91 (7%)	54 (3.8%)
Referred place		
Govt	28 (32.9%)	21 (43.8%)
Private	57 (67.1%)	27 (56.3%)

Investigations	Most recent levels (Mean ± SD)
FBS (N = 520) (in mg/Dl)	155 ± 47
PPBS (N = 483) (in mg/Dl)	186 ± 83
RBS (N = 116) (in mg/Dl)	124 ± 24
SBP (in mm of Hg)	126 ± 36
DBP (in mm of Hg)	88 ± 15

Table 32.3: Non-Communicable Diseases: Recent investigations and measurements details

 Table 32.4:
 Non-Communicable
 Diseases:
 Managing
 routine
 medical
 expenses
 and
 satisfaction
 scores

Routine medicine expenses	DM (N = 1313)	HTN (N = 1539)	
Income	460 (35%)	487 (31.6%)	
Savings	949 (72.3%)	1083 (70.4%)	
Borrowed	26 (2%)	24 (1.5%)	
Selling property / jewelry	9 (0.7%)	8 (0.5%)	
Insurance	5 (0.4%)	8 (0.5%)	
Satisfaction score Median	8 (8, 9)	8 (8, 9)	
Govt			
Satisfaction score Median	8 (8, 9)	8 (8, 9)	
Private			

C.2.7. Subgroup analysis:

1. Classification based on type of house:

Type of house was classified in to pucca and other (semi pucca and Kutcha clubbed together) for subgroup analysis.

Demographic characteristics:

Table 33 provides the comparison of demographic parameters between pucca and non pucca categories. The distribution of education and occupation category was comparable between pucca and non pucca categories. Higher proportion of individual (24.1%) had completed graduate and above in individuals living in Pucca houses compared to non pucca houses (13.9%). Lower proportion (67.4%) of household members showed their BPL card as against 83.7% in non Pucca houses. Socio economic parameters, possession of motor vehicles, were comparable between pucca and non pucca houses. Higher proportion of individuals were holding Arogya Karnataka health insurance in individuals living in non pucca houses. Lifestyle characteristics were comparable between pucca and non pucca houses. Although, the presence of 2 weeks morbidity was comparable between pucca and non pucca houses, higher proportion of individuals (61.3%) sought treatment in health facility among individuals living in non pucca houses. Higher proportion of individuals (70.1%) belonged to pucca houses were availing treatment from private health facility compared to 58.5% of individuals belonged non pucca houses. Mode of transport for the type of houses was comparable. Individuals belonged to non -pucca houses, the preferred time of consultation morning

followed by evening and after noon. Whereas among individuals in pucca houses, the most preferred time of consultation was morning and evening. Time to reach health facility and time taken to consult the doctor was comparable by type of facility and houses. Among individuals living in non pucca houses, higher proportion mentioned income as main source of income form managing their routine medical expenses, whereas among pucca houses, savings was the major source of income for the same. The median cost spent towards investigations and buying medicine were slightly higher for individuals belonged to pucca houses. More percentage of mother (72.2%) lived in non-pucca houses, availed their ANC and PNC care form public health facility with no significant difference for place of Childbirth. The median cost spent towards ANC, Childbirth and PNC care for mothers belonged to pucca were 5000, 20000, 7000 respectively, which were noted to be higher compared to non – pucca houses. Place of routine visit for diabetic and hypertensive care was comparable with approximately equal proportion preferring public and private health facility between members of pucca and non pucca houses. Reported reasons for preference of public and private health facility by pucca and non pucca house members were like overall study group.

Demographic Parameters	Pucca house	Non pucca house (semi pucca and kutcha)
Marital Status (≥18 years)	13954	2954
Currently married	9793 (70.2%)	1907 (64.6%)
Widowed / Separated / Divorced	1410 (10.1%)	470 (15.9%)
Never married	2353 (16.9%)	451 (15.3%)
Unspecified	398 (2.9%)	126 (4.3%)
Education (≥6 years)	16654	3644
Illiterate / Primary / Middle	5027 (30.2%)	1540 (42.3%)
High school	4482 (26.9%)	977 (26.8%)
PUC	2587 (15.5%)	497 (13.6%)
Graduate and above	4015 (24.1%)	506 (13.9%)
Healthy lifestyle	2083 (16.9%)	172 (6.52%)
Health Insurance	N = 13619	N = 3045
AB / AK	2142 (12.1%)	939 (24.2%)
ESI / CGHS	1273 (7.2%)	254 (6.6%)
Private	1484 (8.4%)	96 (2.5%)
Employers	85 (0.5%)	26 (0.7%)
Not available	12718 (71.9%)	2558 (66.1%)

non pucca houses

Morbidity in last 2	Pucca house		Non pucca hous	е	
weeks			_		
Present	1211 (8.3%)		279 (8.2%)		
Availed treatment					
at					
Govt	170 (29.9%)		71 (41.5%)		
Private	399 (70.1%)		100 (58.5%)		
Medicine system					
Allopathy	543 (95.6%)		167 (97.7%)		
Ayurveda	11 (1.9%)		1 (0.6%)		
Others	14 (2.5%)		3 (1.8%)		
	Public	Private	Public	Private	
Manage routine					
medical expenses					
Income	49 (28.8%)	121 (31.8%)	48 (67.6%)	65 (67%)	
Savings	120 (70.6%)	264 (69.5%)	31 (43.7%)	42 (43.3%)	
Borrowed	1 (0.6%)	7 (1.8%)	0 (0%)	2 (2.1%)	
Selling property / jewelry	1 (0.6%)	1 (0.3%)	1 (1.4%)	0 (0%)	
Insurance	1 (0.6%)	0 (0%)	0 (0%)	1 (1%)	
Consultation cost- Rs	100 (50, 200)	200 (120, 300)	190 (112, 300)	150 (100, 250)	
Testing cost- Rs	800(200, 2000)	300(100, 1375)	500(187.5, 812.5)	500(100, 1500)	
Drugs cost -Rs	290 (100, 500)	300 (150, 500)	250 (80, 500)	250 (150, 500)	
Travel cost- Rs	80 (50, 100)	90 (50, 142.5)	100 (50, 200)	100 (50, 200)	

Table 33.2: Comparison of 2-week morbidity characteristics between individuals having pucca and non pucca houses

Table 33.3: Comparison of maternal health characteristics between individuals having pucca	
and non pucca houses	

	ANC		CHILD	CHILDBIRTH		NC
	Pucca N (%)	Non pucca N (%)	Pucca N (%)	Non pucca N (%)	Pucca N (%)	Non pucca N (%)
Place of care	227	54	647	132	229	54
Govt	116 (50.7%)	39 (72.2%)	462 (71.4%)	101 (76.5%)	116 (50.7%)	39 (72.2%)
Private	111 (48.5%)	15 (27.8%)	185 (28.6%)	31 (23.5%)	111 (48.5%)	5 (27.8%)
Type of Childbirt h			N = 450	N = 101		
Normal			226 (50.2%)	63 (62.4%)		

LSCS			223	38 (37.6%)		
and			(49.6%)			
others						
HH	5000	5000	20000	5000	7000	5000
spend	(2000,2000	(2000,1000	(5000,5000	(4125,1500	(2000,2500	(2000,600
Median	0)	0)	0)	0)	0)	0)
(IQR)						

2. Classification based on households belonging to wards with predominant notified slum and non-slum areas; and whether BPL card was shown or not shown at the time of survey:

Subgroup analysis based on households belonging to wards with predominant notified slum and non-slum areas was carried out. Another subgroup analysis was done which was based on the availability of BPL card at the time of survey. Categorization was done based on the households belonging to wards with predominant notified slum and non-slum areas; and the households showing or not showing the BPL card at the time of interview. Results of these two subgroup analyses were also like what was observed in the results by type of houses. Table no. 34 and 35 provides the comparison of households belonging to wards with predominant slum and non-slum areas; and those showing or not showing the BPL card at the time of interview respectively.

 Table 34.1: Comparison of demographic parameters between households belonging to wards with predominant notified slums and non-slum areas

Demographic Parameters	Slum	Non slum
	N = 6222 (28.8%)	N = 15354 (71.2%)
Marital Status (≥18 years)	N = 4865 (78.2%)	N = 11675 (76%)
Currently married	3295 (67.7%)	8393 (71.9%)
Widowed / Separated / Divorced	577 (11.9%)	1302 (11.2%)
Never married	850 (17.5%)	1696 (14.5%)
Unspecified	143 (2.9%)	284 (2.4%)
Education (≥6 years)	N = 5811 (93.4%)	14057 (91.6%)
Illiterate / Primary / Middle	3314 (57%)	8427 (59.9%)
High school	577 (9.9%)	1304 (9.3%)
PUC	1390 (23.9%)	2722 (19.4%)
Graduate and above	530 (9.1%)	1604 (11.4%)
Healthy lifestyle	830 (13.3%)	1425 (9.3%)
Health Insurance		
AB / AK	789 (12.7%)	2292 (14.9%)
ESI / CGHS	538 (8.6%)	989 (6.4%)
Private	738 (11.9%)	842 (5.5%)
Employers	29 (0.5%)	82 (0.5%)
Not available	4128 (66.4%)	11149 (72.6%)

Morbidity in last 2 weeks	Slum [N = 6222 (28.8%)]		Non slum (71.2%)]	[N = 15354]
Present	432 (8.4%)		1058 (8.2%)	
Availed treatment at				
Govt	78 (35.9%)		163 (31.2%)	
Private	139 (64.1%)		360 (68.8%)	
Medicine system				
Allopathy	207 (95.0%)		503 (96.4%)	
Ayurveda	7 (3.2%)		5 (1%)	
Others	3 (1.4%)		14 (2.7%)	
	Public	Private	Public	Private
Managing routine medical expenses				
Income	30 (38.5)	45 (32.4)	67 (41.1)	141 (39.2)
Savings	45 (57.7)	99 (71.2)	106 (65.0)	227 (63.1)
Borrowed	0	1 (0.7)	0	8 (2.2)
Selling property / jewelry	1 (1.3)	1 (0/.7)	1 (0.6)	1 (0.3)
Insurance	2 (2.6)	0	1 (0.6)	2 (0.6)
Consultation cost in Rs	30 (0, 150)	200 (100, 500)	0 (0, 100)	150 (100, 300)
Testing cost in Rs	150 (10, 500)	100 (10, 500)	35 (0, 745)	100 (10, 900)
Drugs cost in Rs	200 (0, 500)	300(100,	100 (0, 350)	250 (150,
		1000)		500)
Cost spent on travel in Rs	85 (20, 100)	100 (50, 195)	50 (0, 100)	60 (41, 107)

Table 34.2: Comparison of 2-week morbidity characteristics between individuals belonging
to wards with predominant notified slums and non-slum areas

 Table 34.3: Comparison of maternal health characteristics between individuals belonging to wards with predominant notified slums and non-slum areas

	ANC		CHILD	CHILDBIRTH		PNC	
	Slum N (%)	Non slum N (%)	Slum N (%)	Non slum N (%)	Slum N (%)	Non slum N (%)	
Place of care							
Govt	33 (54.1%)	122 (55.0%)	65 (48.9%)	239 (56.5%)	64 (50%)	236 (56.3%)	
Private	28 (45.9%)	98 (44.9%)	67 (50.4%)	183 (43.3%)	64 (50%)	182 (43.4%)	
Type of Childbirth							
Normal			73 (55.7%)	216 (51.4%)			
LSCS and others			58 (44.3%)	203 (48.3%)			

HH spend	7000	5000	25000	10000	5000	5000
Median	(5000,	(2000,	(5000,	(5000,	(2000,	(2125,
(IQR)	20000)	20000)	60000)	39500)	25000)	20000)

Table 35.1: Comparison of health insurance between households showing and not showing BPL card at the time of survey

Demographic Parameters	BPL card shown	BPL card not shown
	N = 15170	N = 6397
Health Insurance		
AB / AK	2742 (18.1%)	339 (5.3%)
ESI / CGHS	864 (5.7%)	663 (10.4%)
Private	734 (4.8%)	846 (13.2%)
Employers	45 (0.3%)	66 (1%)
Not available	10785 (70.8%)	4483 (70.1%)

Table 35.2: Comparison of 2-week morbidity	characteristics	households	showing	and not
showing BPL card at the time of survey				

Morbidity in last 2 weeks	BPL card show	wn	BPL card not shown		
	N = 15170		N = 6397		
Present	990 (7.5%)		500 (10.5%)		
Availed treatment at					
Govt	189 (39%)		52 (20.4%)		
Private	296 (61%)		203 (79.6%)		
Medicine system					
Allopathy	470 (97.1%)		240 (94.1%)		
Ayurveda	5 (1%)		7 (2.7%)		
Others	9 (1.8%)		8 (3.1%)		
	Public	Private	Public	Private	
Managing routine medical expenses					
Income	85 (45%)	131 (46.1%)	12 (23.1%)	55 (28.5%)	
Savings	115 (60.8%)	173 (60.9%)	36 (69.2%)	133 (68.9%)	
Borrowed	0 (0%)	3 (1.1%)	1 (1.9%)	6 (3.1%)	
Selling property / jewelry	2 (1.1%) 1 (0.4%)		0 (0%)	0 (0%)	
Insurance	1 (0.5%) 1 (0.4%)		0 (0%)	0 (0%)	
Consultation cost in Rs	150 (50, 250)	150 (100,	150 (100,	225 (150,	
		300)	300)	500)	
Testing cost in Rs	500 (162.5,	300(100,	500 (350,	500 (200,	
	800)	1500)	3500)	1500)	
Drugs cost in Rs	300 (100,	250 (150,	440 (92.5,		
	500)	500)	500)	700)	
Cost spent on travel in Rs	72.5 (50, 100)	80 (50, 145)	100 (50, 167)	100 (50, 200)	

	ANC		CHILDBIRTH		PNC	
	BPL card shown N (%)	BPL card not shown N (%)	BPL card shown N (%)	BPL card not shown N (%)	BPL card shown N (%)	BPL card not shown N (%)
Place of care	205	78	580	199	205	78
Govt	134 (65.4%)	21 (26.9%)	430 (74.1%)	133 (66.8%)	134 (65.4%)	21 (26.9%)
Private	69 (33.7%)	57 (73.1%)	150 (25.9%)	66 (33.2%)	69 (33.7%)	57 (73.1%)
Type of Childbir th			425	126		
Normal			237 (55.8%)	52 (41.3%)		
LSCS and others			187 (44%)	74 (58.7%)		
HH spend Median (IQR)	5000 (2000,1500 0)	10000 (5000,2500 0)	10000 (5000,3550 0)	25000 (5000,6000 0)	5000 (2000,2000 0)	8000 (3000,3000 0)

Table 35.3: Comparison of maternal health characteristics between individuals showing and not showing BPL card at the time of survey

C.3. Perceptions of patients on services received at health facilities

This section presents results of patients perceptions and experiences of services received at both public and private facilities.

Table 36: Socio-demographic characteristics of patients from public and private facilities

Patient sociodemographic characteristics	Public facility (n=92)	Private facility (n=84)*	Test of Signficance
Age (mean±SD) years [Range]	45.6±16.3 [17-81]	39±17 [3-96]	t test=2.62 (p=0.009)
Sex ✓ Male ✓ Female	37 (40%) 55 (60%)	48 (57%) 36 (43%)	Chi-square=5.03 (p=0.025)
Occupation ✓ Govt/private employed ✓ Business / self-employed ✓ Daily wages	19 (21%) 14 (15%) 4 (4%) 10 (5%)	22 (26%) 6 (7%) 6 (7%) 28 (33%)	Chi-square =19.91 (p=0.0005)

 ✓ Unemployed (Retired / not working/student) ✓ Homemaker 	45 (49%)	22 (26%)	
Education in years - Median (Q1,Q3) ✓ Nil ✓ Primary 0-5 th Std ✓ Secondary (6-12 th Std) ✓ Graduation ✓ Postgraduate	10 (7, 12) 16 (17%) 5 (5%) 51 (55%) 16 (17%) 4 (4%)	10 (12%)	Chi-square=6.26 (p=0.18)
Distance from home to health facility (KMs)- Median (Q1,Q3)	1.0 (0.5, 3)	2 (1,5)	
Duration of travel in minutes - Median (Q1,Q3)	10 (5, 15)	10 (5,15)	
*From private facilities patients were not av attempts - 1 patient from 3 facilities; - 2 patients from 1 facility	ailable despite th	iree successive	

2 patients from 1 facility
3 patients from 1 facility

Nearly half of the patients 45 (49%) from public facilities were homemakers while in private facilities 22 (26%) each, were homemakers and public or private employed. Half of all patients from both public and private facilities [51 (55%) and 42 (50%) respectively)] had secondary education, while 20 (24%) of private health patients were graduates (Table 20). The patients from public and private differed significantly (p<0.05) by sociodemographic characteristics such as age (younger in private facilities); sex (more males seeking services at private facilities); occupation (lesser homemakers among those seeking services from private facilities) but not by education level (Table 20).

The commonest health problem for current visit to the health facility was fever (30% and 39%) in both public and private health facilities; pain (25% in both). Few patients returned for follow-up or check of diabetes (11% and 16%) and hypertension (11% and 10%) from public and private health facilities (Figure 10).

The commonest reason for choice of public health facility was free treatment (29%) and good response of health workers 39%, while for private health facilities it was good consultation (54%) and nearby (27%) as seen in Figure 11.







Figure 10: Patient exit interview: Health problem for the current visit to the facility






The good response of health care workers was mentioned by 33% of patients on what they liked most of the facility and 29% mentioned good consultation of the public facilities. While of patients from private health facilities, 20% mentioned good consultation (Figure 12).





Most of the patients in both public and private facilities (>74%) mentioned they had been examined, their BP was checked, they received a prescription, their condition and medications were explained to them at the current visit (Figure 13).





The satisfaction score of private childbirth facilities was higher than that of public facilities, but this was not significantly different. However, patients from public facilities (UPHCs) were significantly more satisfied with services received than with private clinics (p<0.0001) as seen in Table 21. Figure 14 shows that a few (6%) of patients attending private facilities had poor satisfaction levels to services received.

	Overall		Childbirth facility		UPHCs/Clinics	
	Public	Private	Public	Private	Public	Private
	(n=89)	(n=82)	(n=11)	(n=12)	(n=78)	(n=70)
Mean±SD	71.0±10.8	60.6±17.3	69.2±11.8	73.9±11.1	71.3±10.7	58.3±17.2
t test	t=4.75		t=0.98		t=5.58	
[95% CI]	[6.08-14.72]		[-14.63-5.23]		[8.39-17.60]	
(p value)	(p<0.0001)		(p=0.33)		(p<0.0001)	

Table 37: Overall satisfaction on services as reported by patients at exit interview

*3 patients from public and 2 patients from private did not respond to the scale





Figure 14: Patient exit interview: Level of satisfaction for services received at the health facility


C.4. To identify and explain barriers and facilitators to CPHC

The challenges of health workers (Doctors, nurses, lab technicians and pharmacists) from public facilities and those at field (ANMs and ASHAs) were grouped together and presented in Figure 15 and Figure 16. Ratio of HCP with population was reported as the largest challenge (35%), followed by multi-tasking (20%), lack of supplies and meeting targets (19%) were mentioned by HWs of public facilities. While clinical management was reported by 37% of HCPs of private facilities (Figure 15). Among the field level HCPs, management of targets (28%), ratio of HCP and population (23%) and multi-tasking (21%) were the challenges to provide needed services (Figure 16).

Formative Research for CPHC in Mysuru City




Of the community related challenges 23% and 11% of HCPs from public and private facilities mentioned lack of cooperation of people. Lack of acceptance of people to treatment protocols was mentioned by 15% and 26% respectively (Figure 15). While 41% of field HCPs of public facilities mentioned lack of cooperation of community and 21% mentioned it was difficult to mobilize the community (Figure 16)




C.5. To identify design options for strengthening urban primary health care

Suggestions for improvement of services provided

Suggestions to improve services related to health systems were mentioned by HCPs from UPHCs and private facilities included building (39% versus 24%). Drug availability (34%), better lab facilities (30%) and better equipment and facilities (23%) were mentioned by HCPs from UPHCs. On the spot treatment as well as better hygiene was mentioned by 8% of them from UPHCs as community-related suggestions for improvement. While 35% of HCPs primarily doctors mentioned health education for the community (figure 17). Reimbursement, better ratio of HCPs with population and better salary were three health system related suggestions given by 46%, 38% and 35% respectively of field HCPs (Figure 18). More than a quarter of field HCPs (27%) mentioned mobilization of people through key stakeholder involvement as a suggestion to improve services.




Formative Research for CPHC in Mysuru City



- Leadership & governance
- Health financing
- Health workforce
- Essential products and technologies
 - Health service delivery
 - Health information systems
 - Community ownership and partnership

Figure 18: Suggestions given by field HCPs from UPHCs (n=34)

Sixteen percent of patients mentioned that lab equipment needs to be improved; 9% mentioned the need for specialists, 8% suggested inpatient (IP) facilities for emergency from public health facilities to improve services. The same suggestions were given by 4%, 5% and 2% respectively_of patients from private health facilities (Figure 19).



OUT OF POCKET HEALTH EXPENDITURE IN KARNATAKA

In addition to identifying and analyzing barriers and facilitators to comprehensive primary health care and developing the design option for strengthening urban primary health care, out of pocket health expenditure in Karnataka was also analyzed as part of the CPHC study.

Healthcare expenditure results in better provision of health opportunities. It strengthens human capital and improves productivity by contributing to economic performance and it is important to assess the health expenses in every country. The total heath expenditure for India was estimated to be 3.89 % of the country's GDP. The primary goal of the analysis was to draw inferences on the non-communicable disease (NCD) profile in Karnataka's healthcare system using secondary data.

Objectives:

- 1. To describe the health expenditure towards NCDs in Karnataka
- 2. Examine re-imbursement and out of pocket expenses for health by wealth quintiles and other factors.
- 3. Compare health expenses between private and public health facilities.

Data and Variables:

Data was taken from the National Sample Survey Household Consumption Expenditure (Round 75) from July 2017 to June 2018 and representative for the state of Karnataka. Data was obtained from the Ministry of Statistics and Program Implementation, Govt. of India, to measure the population's expenditure for communicable and non-communicable diseases. Data consisted of the usage of both public and private health care, the cost incurred, various medication kinds regardless of how frequently they are used, the cost of treatment, and various ailments covered. This data has a focus on "out of pocket expenditure" as well as access to government-funded health insurance programmes.

Non-communicable diseases were reported among 1.3% of 1217 inpatients and 1.5% of 666 outpatients in Karnataka. Only 0.45% of the 666 outpatients received reimbursement, compared to 7.8% of the 1217 patients who received it.


The last 365 days' value of medical expenses for inpatients include bed charges, doctor and surgeon fees, diagnostic tests, medications, and other medical costs such as attendant costs, physiotherapy, personal medical appliances, blood and oxygen cylinders. The medical costs for outpatients are the same as those for inpatients and include Ayush medicines, doctor and surgeon fees, additional


medications, diagnostic tests, and other medical expenses. The total expense is made up of transportation costs, other non-medical expenses and medical expenditures. Other non-medical expenses include registration fees, food, transportation for others, escort expenses, and lodging fees, if necessary for the patients.

Our study focused on non-communicable illnesses, such as cancer, blood disorders, endocrine, metabolic, nutritional, psychiatric, and neurological conditions. It also examines injuries and conditions of the eye, ear, cardiovascular, respiratory, gastrointestinal, musculoskeletal, and genitourinary systems among inpatients and outpatients. Diabetes, undernutrition, goiter, and other thyroid problems are among the endocrine, metabolic, and nutritional conditions (including obesity). Mental retardation and disorders, headaches, seizures, strokes, hemiplegia, and memory loss are all types of physiological and neurological ailments.


Figure 20: Non communicable diseases among female inpatients

Non-communicable diseases among Females Inpatients





Cardiovascular, gastrointestinal and Musculo skeletal were the most common non communicable disease noted in both male and females' inpatients. Injuries was the commonest cause of hospitalization in male patients.







Figure 23: Non communicable diseases among male outpatients



Endocrine and cardiovascular were the commonest reasons for the OPS visit for both gender.





Figure 24: Source of total expenditure for inpatient admissions


Figure 25: Source of total expenditure for outpatient care




Wealth Quintiles and their Various Source of Finance to pay Total Expenditure- Outpatients

The Total health expenditure exclusive of the amount reimbursed is borne by the household. For both inpatients and outpatients, saving was the most common source of health expenditure. For inpatients, multiple other kinds of sources have been used as borrowing from friends/relatives.


The net out of pocket expenditure of respondents to health care were partitioned into two components- the amount reimbursed (Total reimbursement) and amount not reimbursed (Total expenditure). The distribution of these components by sex of respondent is given in Figures 26 and 27. The average net and total expenditure for inpatients and outpatient visit were higher for males compared to females.



Medical Expenses contribute maximum to the Total Expenditure among both inpatients and outpatients. Other non-medical expenses were reported for inpatients (**Figure 28**). Expenses for travel were substantial for outpatient visits (**Figure 29**).



Further medication costs were majorly for medicines in the in-patients medical expenditure (**Figure 30**) whereas it was for other medicines in the outpatient expenditure (**Figure 31**).






Figure 32: Distribution of inpatient medical expenditure

The distribution of medical expenditure in inpatient and outpatients visits by wealth quintiles is shown in **Figures 33 and 34**. We can observe that medications account for a larger portion of total medical spending in all wealth quintiles. The highest income quintile's outpatients incurred the lowest cost related to Ayush medications. Inpatient diagnostic test and bed costs are nearly identical among lower income quintiles, but doctor and hospital fees are comparably higher in higher income quintiles.






Figure 33: Components of medical expenditure for inpatients by wealth quintiles-





The distribution of NCDs by age for inpatients shows that in the age upto 18years it was majorly injuries. The proportion of cardiovascular diseases increased in the next two age groups (**Figure 35**).




Figure 35: Distribution of NCDs reported by age among inpatients




Figure 36: Distribution of NCDs reported by age among outpatients

 Table 38: Association between the Reimbursed and Non- reimbursed categories with various

 Socio demographics characteristics

Variable	Category	Reimbursement n(%)	Non-Reimbursement n(%)	P- value
Age in years	0-18	5(4.5%)	106(95.5%)	
	19-59	59(7.97%)	682(92.04%)	0.545
	>60	32(8.7%)	333(91.23%)	

Age Vs Reimbursed and Non- reimbursed categories

Patients who were between the ages of 19 and 59 earned the greatest reimbursement rates of any age group, although there was no significant relationship between age and reimbursement (non-reimbursement).

Gender vs Reimbursement Status

Variable	Category	Reimbursement n(%)	Non- Reimbursement n(%)	P- value
Gender	Male	58(8.05%)	662(91.95%)	0.361
	Female	38(7.6%)	459(92.35%)	

Gender was not a factor in determining reimbursement (or non-reimbursement), and more male patients than female patients obtained reimbursement.

Variable	Category	Reimbursemen t n(%)	Non- Reimbursemen t n(%)	P- valu e
Marital Status	Never Married	10(5.18%)	183(94.81%)	0.27
Status	Currency married	70(8.69%)	735(91.30%)	
	Widowed/Divorced/Separate d	16(7.30%)	203(92.69%)	

Marital Status vs Reimbursement Status

Patients who were then married receive higher reimbursement than those who were not married, but there was no association between marital status and reimbursement (non-reimbursement).

Social Group vs Reimbursement

Variable	Category	Reimbursement n(%)	Non- Reimbursement n(%)	P- value
Social Group	Scheduled Tribes / Scheduled Castes	12(5.35%)	229(95.02%)	<0.05
	Other Backward Class	35(5.87%)	561(94.12%)	
	Others	9(2.65%)	331(97.35%)	

Social class was a factor in determining reimbursement, meaning there was a strong correlation between the two groups and more patients from other backward classes obtained compensation than from other social classes.

Variable	Category	Reimbursement n (%)	Non-Reimbursement n (%)	P- value
Religion	Hinduism	78(7.55%)	956(92.45%)	< 0.05
	Islam	11(7.18%)	142(92.81%)	
	Christianity	6(26.08%)	17(73.91%)	
	Sikhism	0(0.0%)	1(100%)	
	Jainism	1(16.67%)	5(83.34%)	

Religion vs Reimbursement

Religion was a determinant of reimbursement, meaning that there was a strong association between the two groups and that mostly patients belonging to Hinduism received more reimbursement.

Education vs Reimbursement

Variable	Category	Reimbursement n (%)	Non Reimbursement n (%)	P- value
Education	Not Literate Literate 1. Without any Schooling 2. Without formal Schooling 3. Primary 4. Secondary 5. Diploma	8(2.13%) 1(14.28%) 0(0%) 23(5.48%) 25(9.29%) 6(16.67%) 33(29.72%)	366(97.86%) 6(85.71%) 1(100%) 396(94.51%) 244(90.70%) 30(83.34%) 78(70.28%)	< 0.05
	6. Graduate			

General education was a factor in reimbursement, meaning that there was a strong association between the two groups and that more educated patients were reimbursed than illiterate patients.

Variable	Category	Reimbursement n(%)	Non-Reimbursement n(%)	P- value
Type of health facility	Govt/Public hospital (incl. HSC/PHC / CHC etc)	1(0.33%)	296(99.67%)	< 0.05
	Private Hospital	93(10.56%)	788(89.45%)	

Type of health facility vs Reimbursement

Type of health facility was also a determinant for Reimbursement (Non-reimbursement) ,that is there was an association between the two groups and Private Hospital inpatients received more reimbursement than any other Medical Institution.

Place of Residence vs Reimbursement

Variable	Category	Reimbursement n(%)	Non-Reimbursement n(%)	P- value
Place of Residence	Rural	21(3.32%)	612(96.68%)	< 0.01
Residence	Urban	75(9.56%)	509(90.43%)	

Place of Residence was again a factor in determining whether or not a person was reimbursed, meaning that there was a correlation between the two groups and more urban residents were reimbursed than rural ones.

Variable	Category	Reimbursement n(%)	Non-Reimbursement n(%)	P- value
Wealth Quintiles	Lowest	6(2.26%)	259(97.73%)	< 0.05
Quintines	Low	7(3.13%)	216(96.86%)	
	Middle	11(4.03%)	262(95.97%)	
	High	21(9.90%)	191(90.09%)	
	Highest	51(20.90%)	193(79.09%)	

Wealth Quintile vs Reimbursement

There was an association between the two categories, that is patients who belong to the highest quintile of wealth received greater compensation, making wealth a determinant of reimbursement (or non-reimbursement.

Employment type vs Reimbursement

Variable	Category	Reimbursement n(%)	Non- Reimbursement n(%)	P- value
Employment type	Self-employed, Casual labour in agriculture	20(3.08%)	628(96.92%)	< 0.05
	Self-employed, Casual labour in non-agriculture	62(17.23%)	298(82.77%)	
	Regular wage/salary earning in agriculture	5(5,32%)	89(94.68%)	
	Regular wage/salary earning in non-agriculture	4(11.76%)	30(88.23%)	
	Others	5(2.47%)	197(97.53%)	

Employment type was a predictor for payment (or non-reimbursement). There was a correlation between the two groups, and patients who are self-employed in non-agricultural work received more compensation.

Variable	Category	Mean Expenditure	t value	P- value
Type of Medical Institution	Public	7729.04	-12.125	< 0.01
mottution	Private	44776.28		

 Table 39: Association of type of Medical Institution and Mean Expenditure

The average expenditure is significantly different in the Public and Private Sectors.

Variable	Category	Mean Expenditure	t value	P- value
Place of Residence	Rural	25478.89	-4.5359	< 0.01
	Urban	46113.24		

The average expenditure is significantly different in Rural and Urban Sectors.

Table 41: Association of Various Components of Medical Expenditure and Type of Medical Institution

Variable		Type of Med Institution	ical	t - value	P- value
		Public	Private		
Components of Medical	Doctors/ Surgeon fee	418.55	8008.09	-11.86	< 0.01
Expenditure	Medicines	2838.57	9683.54	-6.87	< 0.01
	Diagnostics fee	1157.19	4262.91	-7.73	< 0.01
	Bed Charges	274.74	4756.2145	-14.60	< 0.01
	Other Medical expenses	929.75	3185.09	- 4.8008,	< 0.01

The average expenditure of each of the medical components is significantly different in the two different types of Medical Institutes.

Components	Lower	Low	Middle	High	Highest	F- value	P- value
Doctors/ Surgeon fee	4647.543	3938.74	4867.63	6567.66	10353.15	6.60	< 0.01
Medicines	4843.77	5659.35	7227.85	6527.95	9423.43	4.75	< 0.01
Diagnostics fee	2266.13	2112.5	2472.1	2979.67	5869.6	9.22	< 0.01
Bed Charges	2516.04	2292.66	3319.22	3581.07	4884.18	5.28	< 0.01
Other Medical expenses	1644.33	1927.72	1820.12	2608.9	3361.3	1.7832	0.13

 Table 42: Association of Wealth Quintiles and Mean of components contributing to Medical

 Expenditure - Inpatients

Table 43: Association of Wealth Quintiles and Mean of components contributing to MedicalExpenditure – Outpatients

Components	Lowest	Low	Middle	High	Highest	F- value	P- value
Doctors/ Surgeon fee	66.7	241.53	88.76	73.23	6.63	13.11	< 0.01
AYUSH Medicines	74.70	148.08	27.05	118.99	3.7	7.27	< 0.01

Other Medicines	331.75	241.36	483.62	409.62	473.00	6.63	< 0.01
Diagnostics tests	48.85	114.68	39.45	67.09	104.62	3.68	0.005
Other Medical Expenses	57.26	60.5	37.60	13.04	13.28	3.58	0.006

In both Inpatients and Outpatients, at least one Wealth Quintiles mean of various components of Medical Expenditure differs significantly from the other.

Cost of medicines was the major expenditure for both inpatients and outpatients. Health expenditure of individuals in lower quintiles of wealth were mostly not reimbursed and they depended on borrowing money from relatives and other sources.

Conclusions:

The community assessment was a population-based survey carried out in Mysuru city with the objective of profiling the community morbidity status, health care-seeking and costs incurred for selected acute and chronic conditions in urban wards of Mysuru city. A total of 6007 households comprising of 21576 individuals were surveyed from 25 randomly selected wards of Mysuru city. Among 6474 women in the reproductive age group, 100 (1.6%) mothers were currently pregnant and pregnancy in the last 3 years were 563 (8.6%). Among 11978 individuals aged above 30 years, 17.3% of were either diabetes or hypertension.

Health Insurance:

Two third of the households (67.8%) didn't have any insurance coverage. Only 17% of the households were covered under Ayushman Bharat / Aarogya Karnataka. ESI / CGHS and private insurance coverage was 7.8% and 7.4% respectively. Only 26 households utilized their health insurance in the past one year (link it with morbidity profiling)

Morbidity status:

Illness in the last 2 weeks was observed in 7% (n=1490) of the population studied, of which almost half of them sought treatment at health facilities indicating moderate utilization of health facilities for acute illnesses. Among those who sought treatment, only 32.5% received treatment from public health facility. The use of public health care facilities was noted to be low compared to private

health care facilities. Self-medication and use of Over the Counter (OTC) drugs were the reasons reported among those who did not seek treatment at health facility.

Among those who sought treatment at health facilities, 9% reported changing place of treatment after the first visit. Most of them reported that less facilities and long distance were the reasons to change place of treatment. No significant difference between public and private facilities in terms of change of place of treatment was observed. Only 34 individuals reported to have had 2 or more ailments in the last 2 weeks. Distance to health facility, time taken to reach the health facility and time taken to consult the doctor were comparable between individuals seeking care at public and private health facilities, indicating the preference for choosing health facility was not governed by the above said factors. Income and savings were the most used mode for managing their routine medical expenses and were comparable between individuals choosing public and private health facilities. The preference for health facilities was comparable by gender, however, significant difference was seen by age categories. Significantly higher proportion of children between 6-18 years were consulted in private health facilities, which could be because of availability of paediatric specialist in the private set up.

The cost incurred for healthcare in public health facilities was very less as compared to private health facilities. Although there was no / minimal charge of consultation in public health facilities, the median investigation cost and the drug cost was Rs. 65 (10, 520) and Rs. 110 (0, 425) respectively. Patients at exit interview clearly pointed to the need for costs to be affordable, especially in private health facilities, but also in public health facilities since they had to seek diagnostic services outside. This could be due to non-availability of drugs and investigations required in the public health facilities which was highlighted as a suggestion for improving services by HCPs from facilities as well as by patients at exit interviews. But for the private health facilities, cost spent on drugs and investigations was two folds as compared to public health facilities.

Maternal health:

Ante natal care (ANC) was elicited only among the current pregnancies (n = 100) during the study period. All these pregnancies were registered and 46% of them had reported utilizing public health facilities for ANC care. Regarding the cost spent towards ANC care, those who preferred private health facilities had reported to spend five folds of what was spent in public health facilities (Median cost in Pvt = Rs 15,000, Govt. = Rs 3,000)

The data on Childbirth was recorded from the mothers who delivered in the past 3 years. More than half of the mothers utilized public health facilities for their deliveries, and a 53% of them had normal vaginal Childbirth. Significantly higher proportion of females had C-section in private health facilities (67%) compared to only 25% in public health facilities (p<0.01). Similarly for Postnatal

care also, 55% of them preferred public health facilities. Although all UPHCs conducted ANC, the opportunity for childbirth in these settings was non-existent. Patients thus clearly pointed to the need for comprehensive maternity services – ANC, childbirth, and PNC.

For all maternal health services, income and savings were the most common utilized modes for managing medical expenses. Only 1% of them utilized health insurance for their childbirth purpose. The cost spent towards childbirth care was significantly higher among those who utilized private health facilities compared to public health facilities (Median cost in Pvt = Rs 50,000, Govt. = Rs 5,000). PNC care expenses were also noted to be higher in the private as compared to public health facilities.

The reasons for choosing the public health services for maternal health care reported were near distance and free of cost. Good doctor, timely service, and all facilities available at one place were the primary reasons for preferring private health facilities. Like the pattern observed for acute illnesses, the distance, time and cost spent were not determining factors for choosing MCH facilities.

Child Health (≤ 60 months):

About a quarter of the children (25%) were reported to be sick in the last 1 month. Acute Diarrhoeal Disease (ADD) (67.6%) was the most reported illness followed by Acute Respiratory Infection (ARI) (35.5%). For both the ailments, approximately equal proportion of households availed care from public and private health facilities. The median cost spent on treatment for both ARI and ADD in public was one third of what was spent in private health facility. The preferred reasons for choosing public health facility were less / free of cost followed by trust in doctor and nearby distance which was similar for both ARI and ADD. Trust in doctor, timely service, and all facilities at one place were the reasons reported in favour of private health facility. Hospitalization rate for ARI and ADD were 13.2% and 4.0% respectively. Majority of them were hospitalized in private health facilities (ARI – 80% and ADD – 70%). Due to free cost of immunization, majority of children (< 2 years) have been reported to avail child immunization services in the public health facilities.

These findings indicate that although the preference of health facilities for outpatient care was equal in both public and private, but for hospitalization, majority of them preferred private health facilities. This could be due to the availability of comprehensive paediatric care in a private setting. **Non-Communicable Diseases (>30 years):**

The reported prevalence of either diagnosed diabetes or hypertension was 17.3%, (15.9% in males / 18.7% in females). Both diabetes and hypertension were presented in 7.5% of the individuals (6.5% males/ 8.7% females). Diabetes and Hypertension alone was reported in 12.8% and 15.0% respectively.

For NCD care, private health facilities were the preferred health facilities, considering the trust in doctor (80.0%) followed by timely service (50.0%) and all facility at one place (27.0%). Higher proportion of both diabetes (70.3%) and hypertension (65.9%) patients preferred private facility for buying medicine routinely. The regular fasting blood sugar check among diabetic individuals was done in private health facility (69.5%). Even for the NCD complications, most of them were referred to private health facilities.

Non availability of NCD drugs round the year and lack of investigation facilities may be implicated as the reason for inclination towards private health facilities. Like other illnesses, income and savings were reported to be the commonest mode of managing routine medical expenses in NCD patients.

The health facility assessment included walk-through for observation of amenities, infrastructure equipment and supplies of all UPHCs, equivalent number of private health facilities and three public and private childbirth facilities that were less than 30 bedded; interviews with HCPs, and record reviews. In summary, health facilities are easily accessible to the population and located within 1-2 km from the community. The community were able to access services they were seeking within thirty minutes of seeking services. Regular supervision and monitoring of HCPs by a senior within the health facility or health office is occurring. Basic services of ANC, management of minor ailments, first aid for injuries is being managed by UPHCs and private clinics despite HCP shortage. Patients are satisfied with services received and access services based on proximity and their perception of HCPs. This is encouraging despite challenges faced by the HCPs on inadequate facilities, supplies and infrastructure as well as shortage of health workforce. Most of the HCPs at the health facility had received training in relation to common maternal, child, NCD and communicable disease services but only few had received training on RBSK and RKSK which are essential components of services for CPHC. Leadership and governance need to focus towards improving quality of care rather than just quantity. Although majority of HCPs reported to have been supervised or monitored by a senior, this entailed just reviewing reports and targets achieved rather than the quality. Information on financing of the activities at the public and private health facilities was not forthcoming from the senior level HCP.

The facility assessment clearly pointed to gaps in the provision of care due to vital shortage of health workforce coupled with lack of available equipment for diagnostic services, as well as some shortage in supply of essential medications for management of NCDs. The availability of services was limited to 7 hours by all the UPHCs and 5 hours by the private clinics. Only those facilities that provided childbirth services were functional 24/7. Both facility and field HCPs highlighted in the need to improve the building / equipment / lab / maternal services. Services at public health facilities

were mostly accessed by homemakers and women while the private health facilities were accessed mostly by males and younger age group. Health information system is non-existent in both public and private health facilities with no continuity of services for individuals who seek services, especially children and those adults with NCDs. Feedback from the community or individuals is often not obtained to determine ways to improve access, quality, and availability of services.

Out-of-Pocket Health Expenditure: Cost of medicines was the major expenditure for both inpatients and outpatients. Health expenditure of individuals belonging to general category, illiterate, rural residents, and lower quintiles of wealth were mostly not reimbursed and they depended on borrowing money from relatives and other sources. Average expenditure of consultation, drugs and investigation cost were significantly higher for the private health facilities. Also, the average expenditure for inpatients was higher for individuals belonging to urban and higher wealth quintiles.

Organisation approaches that would require improvement includes a more robust health information system that not only facilitates registration of patients so that follow-ups and linkages between facility and field HCPs are planned strategically especially for those with chronic NCDs and CDs but would also aid in monitoring progress with meeting targets. Coupled with workforce shortage both at the health facility and the field, capacity building of HCPs at all levels must be geared towards better communication with patients, identification of complications and appropriate referrals, linkages between public and private health facilities that probably use a common UID for patients to facilitate efficient follow-ups. Capacity building could be facilitated by using the mentoring approach rather than the monitoring and supervision approach. It would be prudent for public health facilities to be re-organised so that there is at least one facility offering childbirth services attached to 4-5 UPHCs. Moreover given the health workforce shortage, a system to make diagnostic services more efficient, yet accessible could include sample collection at the UPHCs with an effort to transport samples to a referral diagnostic centre that would report back to the UPHC details of the test result. This will require a better health information system that links all UPHCs with the referral diagnostic centre.

ICD Mortality Data The distribution of mortality data by ICD classification is provided in table no. 44.

	Male	Female	Total
	n-7692	n-3770	n-11462
Sepsis	319 (4.1)	162 (4.3)	481 (4.2)
Acute Respiratory Infection	83 (1.1)	83 (2.2)	166 (1.4)
HIV/AIDS	12 (0.2)	5 (0.1)	17 (0.1)
Diarrheal Diseases	44 (0.6)	37 (1.0)	81 (0.7)
Meningitis	34 (0.4)	43 (1.1)	77 (0.7)
Pulmonary TB	195 (2.5)	48 (1.3)	243 (2.1)
Haemorrhagic fever	52 (0.7)	50 (1.3)	102 (0.9)
Unspecified Infectious illness	70 (0.9)	41 (1.1)	111 (1.0)
Oral Neoplasm	27 (0.4)	4 (0.1)	31 (0.3)
Unspecified neoplasms	153 (2.0)	66(1.8)	219 (1.9)
Digestive neoplasms	201 (2.6)	79 (2.1)	280 (2.4)
Respiratory neoplasm	79 (1.0)	31 (0.8)	110 (1.0)
Breast neoplasm	0	65 (1.7)	65 (0.6)
Female reproductive neoplasm	0	64 (1.7)	64 (0.6)
Male reproductive neoplasm	30 (0.4)	0	30 (0.3)
Unspecified non communicable disease	1624 (21.1)	625 (16.6)	2249 (19.6)
Diabetes mellitus	557 (7.2)	299 (7.9)	856 (7.5)
Severe malnutrition	18 (0.2)	9 (0.2)	27 (0.2)
Severe anaemia	74 (1.0)	59 (1.6)	133 (1.2)
Unspecified cardiac disease	1151 (15.0)	749 (19.9)	1900 (16.6)
Acute cardiac disease	1043 (13.6)	474 (12.6)	1517 (13.2)

Stroke	464 (6.0)	228 (6.0)	692 (6.0)
COPD	435 (5.7)	92 (2.4)	527 (4.6)
Asthma	17 (0.2)	36 (1.0)	53 (0.5)
Pregnancy related	0	5 (0.1)	5 (0.0)
Prematurity	111 (1.4)	63 (1.7)	174 (1.5)
Birth asphyxia	35 (0.5)	26 (0.7)	61 (0.5)
Neonatal Pneumonia	27 (0.4)	6(0.2	33 (0.3)
Unspecified perinatal cause	2 (0.3)	10 (0.3)	3 (0.3)
Neonatal Sepsis	82 (1.1)	58 (1.5)	140 (1.2)
Congenital formation	76 (1.0)	36 (1.0)	112 (1.0)
Unspecified external cause of death	617 (8.0)	196 (5.2)	813 (7.1)
Epilepsy	37 (0.5)	21 (0.6)	58 (0.5)

Table No. 44: ICD classification by gender

	0-28 days	1 -1 4 years	>14 years
	n-322	n-328	n-10274
Diarrheal Diseases	6 (1.9)	8 (2.4)	65 (0.6)
Pulmonary TB	0	5 (1.5)	230 (2.2)
Unspecified Infectious	0	22 (6.6)	85 (0.8)
Sepsis	0	13 (3.9)	440 (4.3)
Haemorrhagic fever	0	20 (6.0)	78 (0.8)
HIV/AIDS	0	0	17 (0.2)
Oral Neoplasm	0	0	29 (0.3)
Unspecified neoplasms	0	7 (2.1)	208 (2.0)
Digestive neoplasms	0	0	265 (2.6)
Respiratory neoplasm	0	0	107 (1.0)
Breast neoplasm	0	0	62 (0.6)
eoplasm	0	0	61 (0.6)
Male reproductive neoplasm	0	0	30 (0.3)
Severe anaemia	0	2 (0.6)	121 (1.2)
Unspecified non communicable disease	0	22 (6.6)	2095 (20.4)
Diabetes mellitus	0	4(0.3)	797 (7.8)
Severe malnutrition	0	1 (0.3)	20 (0.2)
Meningitis	1 (0.3)	17 (5.1)	58 (0.6)
Unspecified cardiac disease	0	0	1827 (17.8)
Acute cardiac disease	0	0	1471 (14.3)

Stroke	0	0	687 (6.7)
Acute RI	0	3 (0.9)	162(1.6)
COPD	0	3 (0.9)	506 (4.9)
Asthma	0	0	52 (5.9)
Pregnancy related	0	0	5 (0.5)
Prematurity	157 (48.8)	14 (4.2)	0
Birth asphyxia	59 (18.3)	2 (0.6)	0
Neonatal Pneumonia	13 (4.0)	11 (3.3)	0
Unspecified perinatal cause	22 (6.8)	1 (0.3)	0
Neonatal Sepsis	49 (15.2)	54 (16.1)	0
Congenital formation	15 (4.7)	92 (27.5)	0
Unspecified external cause of death	0	24 (7.2)	756 (7.4)
Epilepsy	0	13 (3.9)	40 (0.40

Appendices

Appendix 1: Form 1a: Government - HR Information

- 2. Location /ANM area....
- 3. Date of assessment:
- 4. Name of the investigator:
- 5. Type of facility:
 - i. UCHC...iiUPHC ...iii.ESI Dispensary...ivHealth Kiosk...vother (specify)

ii. Complete the following information

	Doctor	Staff	ANM	ASHA	Lab	Pharmacist	DEO	Class D
		Nurse			technician			
Number of								
sanctioned posts								
Number of posts								
currently filled								
Number of posts								
currently								
available								

iii. Number of sanctioned staff posts not filled

S.	Staff position not filled	Reason	Duration the post
No	1. Doctor	1. No one coming for post	vacant
	2. Staff nurse	2. Not recruited	1. <6 months
	3. ANM	3. Other specify	2. 6-12 months
	4. ASHA		3. >12 months
	5. Pharmacist		
	6. Lab technician		
	7. DEO		
	8. Class D		
1.			
2.			

iv. Record the name of staff available currently and complete the details

S.No.	Name of	Ge	nder	De	signation	Educational	Ap	pointment of	Ho	urs at work
	the staff	1.	Male	1.	МО	Qualification:	the	staff	1.	7am-1pm
	currently	2.	Female	2.	Doctor	1. MD / MS	1.	Permanent	2.	4pm-8pm
	available	3.	Other	3.	Nurse	specialist		/regular at	3.	Otherspecify
				4.	ANM	2. MBBS		current		
				5.	Pharmacist	3. AYUSH		health		
				6.	Pharmacy	4. GNM		facility		
					Asst	5. BSc (Nsg)	2.	On		
				7.	Lab	6. B. Pharm		deputation		
					technician	7. Dip Pharm		from another		
				8.	ASHA	8. DMLT/		health		
				9.	Otherspecify	BSc MLT		facility		
						9. Otherspecify	3.	Shared with		
								another		
								facility		
							4.	Contracted		
								full time		
							5.	Contracted		
								part time		
1.										
2.										
3.										
4.										
5.										
6.										
7.										
8.										
9.										
10.										
11.										

S.No.	Name of	Gender	Designation	Educational	Reason for not	Duration
	the staff	1. Male	1. MO	Qualification:	being available:	staff not
	currently	2. Female	2. Doctor	1. MD / MS	1. On	available
	not	3. Other	3. Nurse	specialist	deputation	1. <6
	available		4. ANM	2. MBBS	to another	months
			5. Pharmacist	3. AYUSH	facility	2. 6-12
			6. Pharmacy	4. GNM	2. On leave /	months
			Asst	5. BSc (Nsg)	pursuing	3. >12
			7. Lab	6. B. Pharm	higher	months
			technician	7. Dip Pharm	education	
			8. Other	8. DMLT/ BSc	3. Not	
			specify	MLT	reporting for	
				9. Other specify	duty	
					4. Other	
					specify	
1.						
2.						
3.						
4.						

v. Number of regular staff posts currently not available

Appendix 2: Form 1b: Private - HR Information

- 1. Name of the Facility:
- 2. Location /ANM area.....
- 3. Date of assessment:
- 4. Name of the investigator:
- 5. Type of facility:
 - B. Private
 - i. Private/Corporate Allopathy Hospital
 - ii. Private/Corporate Allopathy Clinic
 - iii. Private/Corporate AYUSH Hospital
 - iv. Private/Corporate AYUSH Clinic
 - v. NGO/FBO Allopathy Hospital
 - vi. NGO/FBO Allopathy clinic
 - vii. RMP
 - viii. Other (specify)

6. Record the name of staff available currently and complete the details

S. No.	Name of the staff currently available	Gender 1. Male 2. Female 3. Other	 Designation 1. MO 2. Doctor 3. Nurse 4. ANM 5. Pharmacist 6. Pharmacy Asst 7. Lab technician 8. Other specify 	Educational Qualification: 1. MD / MS specialist 2. MBBS 3. AYUSH 4. GNM 5. BSc (Nsg) 6. B. Pharm 7. Dip Pharm 8. DMLT/ BSc MLT 9. Other specify	 Appointment of the staff Permanent /regular at current health facility Contracted full time Contracted part time Other specify 	Hours of work1. morning2. evening3. Other specify
1						
2						

Appendix 3: Form 2a: Govt – staff interview

Instructions: Complete the information in Section A. Section A-G would provide information of the Health Care facility services, timings, challenges faced.

- Section B provides information from Doctors. Select one doctor in the facility.
- Section C provides information of the from Nurses. Select one nurse in the facility
- Section D provides information from Pharmacist. Get him/her to complete the form
- Section E provides information from the Lab Technician. Get him/her to complete the form
- Section F provides information from the ANM. Get him/her to complete the form
- Section G provides information from the ASHA. Get him/her to complete the form

Section A

- 1. Name of the Facility:
- 2. Type of facility:
 - i. UCHC
 - ii. UPHC
- iii. ESI dispensary
- iv. Health Kiosks
- v. Other (specify)
- 3. Location & ANM area:
- 4. Date of assessment:
- 5. Name of the investigator:

SECTION B (For Doctor)

Baseline information

- 1. Experience (years):
- 2. Did you attend any Training / Workshop / Conference in the last 5 years on the following topics (*Tick all those that are applicable*)?
 - a. Population health disease burden assessment and health planning (for doctors only)
 - b. Skilled birth attendance: Specify.....

- c. Newborn care
 - i. Basic newborn care
 - ii. Resuscitation
 - iii. Kangaroo Mother Care for LBW babies
 - iv. Care of small babies Facility based newborn care
- d. RashtriyaBalSwasthyaKaryakram(RBSK) [Identification and early intervention on any of the four Ds for children 1-18 years]
 - i. Defects at birth
 - ii. Deficiencies
 - iii. Diseases
 - iv. Development delays and disability
- e. Infection Control
- f. Family planning
- g. RashtriyaKishorSwasthyaKaryakram or Adolescent Health
- h. Non communicable diseases
 - i. Diabetes
 - ii. Hypertension
 - iii. Cancers
 - iv. Injuries
 - v. Mental Health and wellness
- i. Communicable diseases
 - i. Tuberculosis
 - ii. Dengue
 - iii. Diarrhoea
 - iv. Other communicable diseases
- 3. Age (years):
- 4. Population Area of responsibilities.....
- 5. What services are provided in this facility? (*Tick all services relevant in box provided/timings when services are provided/days when services are provided*)

Maternity	Child health	General Adult	NCD (Diabetes/	Communicable
			HTN)	(TB)

	ANC		Immunisation		Treatment of		Check-ups		DOTS
	Immunisation		Treatment of minor		minor illness		Follow-up		Follow-up
	IFA / Ca		illness / problems		Medical		Eye check		
	Contraception		Growth monitoring		certificated		Kidney		
			First aid for injuries		First aid for		Check		
					minor injuries		ECG		
							Basic drugs		
Tiı	me:	Tiı	ne:	Ti	me:	Ti	me:	Tiı	ne:
	7am-1pm		7am-1pm		7am-1pm		7am-1pm		7am-1pm
	4pm-8pm		4pm-8pm		4pm-8pm		4pm-8pm		4pm-8pm
	24/7		24/7		24/7		24/7		24/7
Da	lys	Da	ys	Da	ys	Da	iys	Da	ys
Da	ys Monday	Da	ys Monday	Da	ys Monday	Da	ys Monday	Da	ys Monday
					·		•		•
	Monday		Monday		Monday		Monday		Monday
	Monday Tuesday		Monday Tuesday		Monday Tuesday		Monday Tuesday		Monday Tuesday
	Monday Tuesday Wednesday		Monday Tuesday Wednesday		Monday Tuesday Wednesday		Monday Tuesday Wednesday		Monday Tuesday Wednesday
	Monday Tuesday Wednesday Thursday		Monday Tuesday Wednesday Thursday		Monday Tuesday Wednesday Thursday		Monday Tuesday Wednesday Thursday		Monday Tuesday Wednesday Thursday
	Monday Tuesday Wednesday Thursday Friday		Monday Tuesday Wednesday Thursday Friday		Monday Tuesday Wednesday Thursday Friday		Monday Tuesday Wednesday Thursday Friday		Monday Tuesday Wednesday Thursday Friday
	Monday Tuesday Wednesday Thursday Friday Saturday		Monday Tuesday Wednesday Thursday Friday Saturday		Monday Tuesday Wednesday Thursday Friday Saturday		Monday Tuesday Wednesday Thursday Friday Saturday		Monday Tuesday Wednesday Thursday Friday Saturday
	Monday Tuesday Wednesday Thursday Friday Saturday Sunday		Monday Tuesday Wednesday Thursday Friday Saturday Sunday		Monday Tuesday Wednesday Thursday Friday Saturday Sunday		Monday Tuesday Wednesday Thursday Friday Saturday Sunday		Monday Tuesday Wednesday Thursday Friday Saturday Sunday

- 6 Did you have supervision / support from a senior person in the last 3 months?
 - □ Yes
 - 🗆 No
- 7 What are the challenges you face in providing the required services?
 - □ Lack of facilities/supplies/equipments/drugs
 - □ Doctor/supervision not available all the time
 - □ Poor linkage with other labs/hospitals/specialists/community
 - □ Lack of staff/high turnover/over worked/less salary
 - □ Lack of training opportunities/promotion /support
 - □ Others (specify).....

SECTION C (For Nurses)

Baseline information

- 1. Experience (years):
- 2. Did you attend any Training / Workshop / Conference in the last 5 years on the following topics (*Tick all those that are applicable*)?
 - a. Skilled birth attendance: Specify.....
 - b. Newborn care
 - i. Basic newborn care
 - ii. Resuscitation
 - iii. Kangaroo Mother Care for LBW babies
 - iv. Care of small babies Facility based newborn care
 - c. RashtriyaBalSwasthyaKaryakram (Identification and early intervention on any of the

four Ds for children 1-18 years)

- i. Defects at birth
- ii. Deficiencies
- iii. Diseases
- iv. Development delays and disability
- d. Infection Control
- e. Family planning

- f. RashtriyaKishorSwasthyaKaryakram or on Adolescent Health
- g. Non communicable diseases
 - i. Diabetes
 - ii. Hypertension
 - iii. Cancers
 - iv. Injuries
 - v. Mental Health and wellness
- h. Communicable diseases
 - i. Tuberculosis
 - ii. Dengue
 - iii. Diarrhoea
 - iv. Other communicable diseases
- 3. Age (years):
- 4. What are the services you are providing in the health facility?

Ma	aternity	Ch	ild health	Ge	neral Adult	NCD (Diabetes/ HTN)		Communicable	
								(TI	B)
	ANC		Immunisation		Check vitals		Check vitals		DOTS
	registration		Injections		Explain		Follow-up		Follow-up
	Immunisation		Growth		medications		Remind for Eye		reminder
	IFA / Ca		monitoring		Dressing		check		Sputum
	Insertion of		Dressings		Injections		Blood sampling		check
	CuT		Splinting for		Starting IV		Do ECG		
	Oral		fracture		Taking ECG		Check foot		
	contraceptives		ORS		Blood		Explain on		
					sampling		physical activity		
							Others		
Tir	me:	Tiı	ne:	Tiı	ne:	Tiı	me:	Tiı	me:
	7am-1pm		7am-1pm		7am-1pm		7am-1pm		7am-1pm
	4pm-8pm		4pm-8pm		4pm-8pm		4pm-8pm		4pm-8pm
	24/7		24/7		24/7		24/7		24/7
Da	ys	Da	ys	Da	ys	Da	ays	Da	ys
	Monday		Monday		Monday		Monday		Monday
	Tuesday		Tuesday		Tuesday		Tuesday		Tuesday
	Wednesday		Wednesday		Wednesday		Wednesday		Wednesday
	Thursday		Thursday		Thursday		Thursday		Thursday
	Friday		Friday		Friday		Friday		Friday
	Saturday		Saturday		Saturday		Saturday		Saturday
	Sunday		Sunday		Sunday		Sunday		Sunday
	All days		All days except		All days		All days except		All days
	except Sunday		Sunday		except Sunday		Sunday		except
									Sunday

- 5. Did you have supervision / support from a senior person in the last 3 months?
 - □ Yes
 - □ No

- 6. What are the challenges you face in providing the required services?
 - □ Lack of facilities/supplies/equipments/drugs
 - □ Doctor/supervision not available all the time
 - □ Poor linkage with other labs/hospitals/specialists/community
 - □ Lack of staff/high turnover/over worked/less salary/more responsibilities
 - □ Lack of training opportunities/promotion /support
 - □ Others (specify).....

SECTION D (For Lab technician)

Baseline information

- 1 Experience (years):
- 2 Age (years):
- 3 Type of employment: Outsourced Employee
- 4 Did you attend any Training / Workshop / Conference in the last 5 years Specify details (Topic)
- 5. What Lab services are provided in this facility?

Maternity	General Child/Adult
🗆 Hb	🗆 Hb
□ Hepatitis	
🗆 Malaria	\square PPBS
□ VDRL	□ FBS
□ Urine analysis	□ Hb1Ac
	Renal Function Test

	Dengue
	□ TB-Sputum
	□ other
Time:	Time:
□ 7am-1pm	□ 7am-1pm
□ 4pm-8pm	□ 4pm-8pm
□ 24/7	□ 24/7
Days	Days
□ Monday	□ Monday
	-
□ Tuesday	□ Tuesday
-	TuesdayWednesday
□ Tuesday	, i i i i i i i i i i i i i i i i i i i
TuesdayWednesday	□ Wednesday
 Tuesday Wednesday Thursday 	WednesdayThursday
 Tuesday Wednesday Thursday Friday 	 Wednesday Thursday Friday
 Tuesday Wednesday Thursday Friday Saturday 	 Wednesday Thursday Friday Saturday

- 6. Do you have linkage with a referral lab? Yes No
- 7. Did you have any kind of support/supervision by officials in the last 3 months at laboratory?
 - □ Yes
 - □ No
- 8. What are the challenges you face in providing the required services?
 - □ Lack of facilities/supplies/equipments/drugs
 - □ Doctor/supervision not available all the time
 - □ Poor linkage with other labs/hospitals/specialists/community
 - □ Lack of staff/high turnover/over worked/less salary
 - □ Lack of training opportunities/promotion /support
 - □ Others (specify).....

SECTION E (For Pharmacist)

Baseline information

- 1 Experience (years):
- 2 Age (years):
- 3 Type of employment: Outsourced Employee
- 4 Did you attend any Training / Workshop / Conference in the last 5 years Specify details (Topic)

5 What is the time period when the services are provided? (*Tick which days services are provided against each option*

Time: to (Morning)

Mon	Tues Wed	Thurs Friday	Sat Sunday
Time:		to	(Evening) / All days / only
Mon	Tues Wed	Thurs Friday	Sat Sunday

Day and night / 24/7

Mon Tues Wed Thurs Friday Sat Sunday

- 6 Did you have any kind of support/supervision by officials in the last 3 months at pharmacy?
 - □ Yes
 - □ No
- 7 What are the challenges you face in providing the required services?
 - □ Lack of facilities/supplies/equipments/drugs
 - □ Doctor/supervision not available all the time
 - □ Poor linkage with other labs/hospitals/specialists/community
 - □ Lack of staff/high turnover/over worked/less salary

- □ Lack of training opportunities/promotion /support
- □ Others (specify).....

SECTION F (For ANM)

Baseline information

- 1 Experience (years):
- 2 Did you attend any Training / Workshop / Conference in the last 5 years on the following topics (*Tick all those that are applicable*)?
 - a. Skilled birth attendance: Specify.....
 - b. Newborn care
 - i. Basic newborn care
 - ii. Resuscitation
 - iii. Kangaroo Mother Care for LBW babies
 - iv. Care of small babies HBNC
 - c. RashtriyaBalSwasthyaKaryakram (Identification and early intervention on any of the four Ds for children 1-18 years)
 - i. Defects at birth
 - ii. Deficiencies
 - iii. Diseases
 - iv. Development delays and disability
 - d. Infection Control
 - e. Family planning
 - f. RashtriyaKishorSwasthyaKaryakram or on Adolescent Health
 - g. Non communicable diseases
 - i. Diabetes
 - ii. Hypertension
 - iii. Cancers
 - iv. Injuries
 - v. Mental Health and wellness
 - h. Communicable diseases
 - i. Tuberculosis
 - ii. Dengue
 - iii. Diarrhoea

iv. Other communicable diseases

- 3 Age (years):
- 4 What are the outreach services you are providing for the health facility?

Ma	iternity	Ch	ild health	Ge	neral Adult	NC	CD (Diabetes/	Co	mmunicable
						НЛ	TN)	(T]	B)
	ANC		Immunisation		Check vitals		Check vitals		DOTS
	registration		Growth		Explain		Follow-up		Follow-up
	Immunisation		monitoring		medications		Remind for		reminder
	IFA / Ca		Dressings		Dressing		Eye check		Remind for
	Oral		ORS				Check foot		sputum
	contraceptives						Explain on		check
							physical		
							activity		
Tiı	ne:	Ti	me:	Ti	me:	Ti	me:	Ti	me:
	7am-1pm		7am-1pm		7am-1pm		7am-1pm		7am-1pm
	4pm-8pm		4pm-8pm		4pm-8pm		4pm-8pm		4pm-8pm
	24/7		24/7		24/7		24/7		24/7
Da	ys	Da	lys	Da	ys	Da	ys	Da	ys
	Monday		Monday		Monday		Monday		Monday
	Tuesday		Tuesday		Tuesday		Tuesday		Tuesday
	Wednesday		Wednesday		Wednesday		Wednesday		Wednesday
	Thursday		Thursday		Thursday		Thursday		Thursday
	Friday		Friday		Friday		Friday		Friday
	Saturday		Saturday		Saturday		Saturday		Saturday
	Sunday		Sunday		Sunday		Sunday		Sunday
	All days		All days		All days		All days		All days
	except Sunday		except Sunday		except		except		except
					Sunday		Sunday		Sunday

- 5 Did you have any kind of support/supervision by officials in the last 3 months?
 - □ Yes
 - □ No
- 6 What are the challenges you face in providing the required services?
 - □ Lack of facilities/supplies/equipments/drugs
 - □ Doctor/supervision not available all the time
 - □ Poor linkage with other labs/hospitals/specialists/community
 - □ Lack of staff/high turnover/over worked/less salary
 - □ Lack of training opportunities/promotion /support
 - □ Others (specify).....

SECTION G (For ASHA)

Baseline information

- 1. Experience (years):
- 2. Did you attend any Training / Workshop / Conference in the last 5 years on the following topics (*Tick all those that are applicable*)?
 - a. Identification of pregnancies/risk pregnancies/ referrals/planning for deliveries
 - b. Newborn care
 - i. Basic newborn care
 - ii. Kangaroo Mother Care for LBW babies
 - iii. Care of small babies HBNC
 - c. Infection Control
 - d. Family planning
 - e. RashtriyaKishorSwasthyaKaryakram or on Adolescent Health
 - f. Non communicable diseases
 - i. Diabetes
 - ii. Hypertension
 - g. Communicable diseases
 - i. Tuberculosis
 - ii. Dengue
 - iii. Diarrhoea

Formative Research for CPHC in Mysuru City

iv. Other communicable diseases

3. Age (years):

4. What are the outreach services you are providing for the health facility?

Maternity	Child health	NCD (Diabetes/ HTN)	Communicable
			(TB)
	Growth monitoring	□ Call for Follow-up	□ Follow-up
registration	□ ORS	□ Eye check	reminder
		□ Check foot	□ Remind for
		□ Explain on	sputum check
		physical activity	
Time:	Time:	Time:	Time:
□ 7am-1pm	□ 7am-1pm	□ 7am-1pm	□ 7am-1pm
□ 4pm-8pm	□ 4pm-8pm	□ 4pm-8pm	□ 4pm-8pm
□ 24/7	□ 24/7	□ 24/7	□ 24/7
Days	Days	Days	Days
□ Monday	□ Monday	□ Monday	□ Monday
□ Tuesday		□ Tuesday	□ Tuesday
□ Wednesday	□ Wednesday	□ Wednesday	□ Wednesday
□ Thursday	□ Thursday		□ Thursday
□ Friday	□ Friday	□ Friday	🗆 Friday
□ Saturday	□ Saturday	□ Saturday	□ Saturday
	□ Sunday		
□ All days	□ All days except	□ All days except	□ All days except
except Sunday	Sunday	Sunday	Sunday

- 5. Did you have supervision / support for a field visit from a senior person in the last 3 months?
 - □ Yes
 - □ No
- 6. What are the challenges you face in providing the required services?

- □ Lack of facilities/supplies/equipments/drugs
- □ Doctor/supervision not available all the time
- □ Poor linkage with other labs/hospitals/specialists/community
- □ Lack of staff/high turnover/over worked/less salary
- □ Lack of training opportunities/promotion /support
- □ Others (specify).....

Appendix 4: Form 2b: Private - staff interview form

Instructions: Complete the information in Section A. Section A-D would provide information of the Health Care facility services, timings, challenges faced.

- Section B provides information from Doctors. Select one doctor in the facility.
- Section C provides information from Nurses. Select one nurse in the facility

Section A

- 1. Name of the Facility:
- 2. Type of facility:
 - vi. Private/Corporate Allopathy Hospital
 - vii. Private/Corporate Allopathy Clinic
 - viii. Private/Corporate AYUSH Hospital
 - ix. Private/Corporate AYUSH Clinic
 - x. NGO/FBO Allopathy Hospital
 - xi. NGO/FBO Allopathy clinic
 - xii. RMP
 - xiii. Other (specify)
- 3. Location:
- 4. Date of assessment:
- 5. Name of the investigator:

SECTION B (For Doctor)

Baseline information

- 1. Experience (years):
- 2. Did you attend any Training / Workshop / Conference in the last 5 years on the following topics -Specify details (Topic)
- 3. Age (years):
- 4. What services are provided in this facility? Please mention during what time period of the day are the services provided (*Tick all services relevant in box provided/timings when services are provided/days when services are provided*)

Ma	iternity	Ch	ild health	Ge	neral Adult	NC	CD (Diabetes/	Co	mmunicable
						ΗΊ	TN)	(TI	3)
	ANC		Immunisation		Treatment of		Check ups		DOTS
	Immunisation		Treatment of		minor illness		Follow-up		Follow-up
	IFA / Ca		minor illness /		Medical		Eye check		
	Contraception		problems		certificates		Kidney		
			Growth		First aid for		ECG		
			monitoring		minor injuries		Basic drugs		
			First aid for						
			injuries						
Tiı	me:	Tiı	me:	Ti	me:	Ti	me:	Tiı	ne:
	morning		morning		morning		morning		morning
	4pm-8pm		4pm-8pm		4pm-8pm		4pm-8pm		4pm-8pm
	24/7		24/7		24/7		24/7		24/7
Da	ys	Da	ys	Da	ys	Da	ys	Da	ys
	Monday		Monday		Monday		Monday		Monday
	Tuesday		Tuesday		Tuesday		Tuesday		Tuesday
	Wednesday		Wednesday		Wednesday		Wednesday		Wednesday
	Thursday		Thursday		Thursday		Thursday		Thursday
	Friday		Friday		Friday		Friday		Friday
	Saturday		Saturday		Saturday		Saturday		Saturday
	Sunday		Sunday		Sunday		Sunday		Sunday
	All days		All days		All days		All days		All days
	except Sunday		except Sunday		except Sunday		except		except
1							Sunday		Sunday

- 5. What are the challenges you face in providing the required services?
 - □ Lack of facilities/supplies/equipments/drugs
 - □ Doctor/supervision not available all the time
 - □ Poor linkage with other labs/hospitals/specialists/community
 - □ Lack of staff/high turnover/over worked/less salary

- □ Lack of training opportunities/promotion /support
- □ Others (specify).....

SECTION C (For Nurse)

Baseline information

- 1 Experience (years):
- 2 Did you attend any Training / Workshop / Conference in the last 5 years on the following topics -Specify details (Topic)
- 3 Age (years):
- 4 What are the services you are providing in the health facility

Maternity	Child health	General Adult	NCD (Diabetes/	Communicable	
			HTN)	(TB)	
□ ANC	□ Immunisation	□ Treatment	□ Check ups	□ DOTS	
□ Immunisation	□ Treatment of	of minor	□ Follow-up	□ Follow-up	
🗆 IFA / Ca	minor illness /	illness	□ Eye check		
□ Contraception	problems	□ First aid	□ Kidney		
	□ Growth	for minor	□ ECG		
	monitoring	injuries	□ Basic drugs		
	□ First aid for				
	injuries				
Time:	Time:	Time:	Time:	Time:	
□ 7am-1pm	□ 7am-1pm	□ 7am-1pm	□ 7am-1pm	□ 7am-1pm	
□ 4pm-8pm	□ 4pm-8pm	□ 4pm-8pm	□ 4pm-8pm	□ 4pm-8pm	
□ 24/7	□ 24/7	□ 24/7	□ 24/7	□ 24/7	
Days	Days	Days	Days	Days	
□ Monday	□ Monday	□ Monday	□ Monday	□ Monday	
□ Tuesday	□ Tuesday	🗆 Tuesday	□ Tuesday	□ Tuesday	
□ Wednesday	□ Wednesday	🗆 Wednesda	□ Wednesday	□ Wednesday	
□ Thursday	□ Thursday	у	□ Thursday		
□ Friday	□ Friday	□ Thursday	□ Friday	🗆 Friday	

Saturday	Saturday	Friday	Saturday	Saturday
Sunday	Sunday	Saturday	Sunday	Sunday
All days	All days	Sunday	All days	All days
except Sunday	except Sunday	All days	except	except
		except	Sunday	Sunday
		Sunday		

- 5 What are the challenges you face in providing the required services?
 - □ Lack of facilities/supplies/equipments/drugs
 - □ Doctor/supervision not available all the time
 - □ Poor linkage with other labs/hospitals/specialists/community
 - □ Lack of staff/high turnover/over worked/less salary
 - □ Lack of training opportunities/promotion /support
 - □ Others (specify).....

Appendix 5: Form 3: Observation Checklist

2.	Location /ANM area
Ζ.	Location / Anim area
3.	Date of assessment:
4.	Name of the investigator:
5.	Type of facility:
	A. Government
	i. UCHC
	ii. UPHC
	iii. ESI Dispensary
	iv. Health Kiosk
	B. Private
	i. Private/Corporate Allopathy Hospital
	ii. Private/Corporate Allopathy Clinic
	iii. Private/Corporate AYUSH Hospital
	iv. Private/Corporate AYUSH Clinic
	v. NGO/FBO Allopathy Hospital
	vi. NGO/FBO Allopathy clinic
	vii. RMP
	viii. Other (specify)
6.	Infrastructure: (Tick Y if available)

6.	Infrastructure: (Tr	ck Y if av	ailable)	
	Functional registration counter	Y	Ν	Not applicable
	Dedicated room for OPD consultation	Y	Ν	
	Examination area in clinic	Y	Ν	
	Labour room	Y	Ν	
	Newborn Corner	Y	Ν	
	Injection room/Dressing room	Y	Ν	
	Lab	Y	Ν	
	Pharmacy	Y	Ν	
	Data entry / records section	Y	Ν	
	Biomedical waste management system	Y	Ν	
	Other facilities i. Clean and functional toilet	Y	Ν	
	ii. Drinking water	Y	Ν	
	iii. Waiting area	Y	Ν	
	iv. Disposal of waste area	Y	Ν	
	v. CSSD/Laundry			
	vi. Easily accessible footpath/ entrance to facility for all patients	Y	Ν	

	vii. Yoga/exercise/meditation hall	Y	N		
	viii. Cooking demonstration area	Y	N		
	ix. Separate room for counseling	Y	N		
	x. Nearest park for exercise	kms	11		
	<u>^</u>				
-		kms			
7.	Equipment and I		<u></u>		1' 1 1
1.	Weighing machine	Y	N	Not A	pplicable
<u>2.</u> 3.	BP Apparatus Stethoscope	Y Y	N N		
<u> </u>	Thermometer	Y	N N		
<u>4.</u> 5.	ECG machine	Y	N N		
<u> </u>	Pulse oximeter	Y	N		
7.	Snellen's Chart	Y	N		
8.	Ophthalmoscope	Y	N		
9.	Dressing kit	Y	N		
).	Thayi card is available and updated(Randomly	EDD	11	Yes /	No
	check any 3 cards)	Weight			110
		BP			
		Bloodgroup	p typing		
		Hb			
		HIV		-	
		Inj.TT			
		Tab.IFA			
		Tab. Calciu	ım		
8.	Lab vis	sit			
1.	Time for blood collection and delivery of reports	Y	N		
	(within 24hrs) displayed				
2.	Availability of drugs and consumables for lab	Y	Ν		
	services				
	i. Stains				
	ii. Reagentsiii. Processing chemicals				
	iv. Rapid diagnostic kits				
	v. Glassware				
	v. Glusswale				
3.	Availability and functionality of equipment and	1			1
	instruments for lab services?				
	i. Hemoglobin meter	Available	Func	tional	
	ii. Differential blood cell counter	Available	Fund	ctional	
	iii. Centrifuge	Available	Fund	ctional	
	iv. Colorimeter	Available		ctional	
	v. Microscope (Malaria, RNTCP/NTEP)	Available	Func	tional	
	vi. Refrigerator	Available		tional	
	vii. Biochemistry analyzer	Available	Func		

4.	Quality control practices a. Calibration of instruments b. Accreditation process		ř ř	N N	Not Applicable
5.	 What are the lab services available at facility? I. ANC kit Pregnancy test, Hb Blood gp and typing TSH HIV test Blood sugar HBsAg VDRL II. Smear test AFB sputum III. Urine tests Urine protein Urine micro IV. Test for Diabetes Blood sugar-FBS, Blood sugar-PPBS HbA1C Lipid profile Serum Creatinine Serum Potassium 	Y N			Not Applicable
6. 7.	Records maintained at laboratory Stock register Indent register Expiry register Test done Comments	Y	N		Not Applicable
9.	Pharmacy	visit			
		Y	N	Not	Applicable
1.	Availability of drugsI.Maternal drugsa.IFAb.Inj.dexamethasonec.AlbendazoleII.Emergency Drugsa.Inj.Adrenalineb.Inj.Hydrocortisonec.Inj.CalciumgluconateIII.Antibiotic drugsIV.VaccinesV.Drugs for TBVI.Drug for MalariaVII.Tobacco cessation drugs (varenicline,		1		Applicable

	NDT ata)				
	VIII. Medicines	used in Diabetes Mellitus			
		fonylureas (glimepiride,			
		pizide, glibenclamide, etc)			
		guanides (metformin)			
		ulin			
	-	pirin/clopidogrel			
	e. Sta	tins (simvastatin, atorvastatin,			
	etc	·			
	IX. Medicines	used for HTN			
	a. AC	E-inhibitor			
	b. AR	B			
	c. Bet	ta-blocker			
	d. Cal	lcium-channel blocker			
	e. Diu	ıretic			
2.		lace to maintain temperature	Y	N	
	chart of deep freeze				
3.	Power backup avail		Y	N	
4.	<u> </u>	ls maintained at pharmacy?	-		
т.	Stock register	is maintained at pharmacy.	Y	Ν	
	Indent register		Y	N	
	Expiry register		Y	N	
10.		Display of n		1	
10.		Display of h	llatti lai		
1.	IEC for	Specify			
	□ Mothers	1 5			
	\Box Adults				
2.	Other displays	Phone Numbers of Referral H	lospitals		
	o unor ansprays	Patient Rights			
		Ambulance contact number			
		KPME (Karnataka Private Me	edical Establis	hment)	
		Registration		linent)	
	Comments:	Registration			
	Comments.				
11.		Infection c	ontrol		
11.	Detiont core cros (I	abour room/ward/Newborn			
1.					
		d equipment are clean.	v	NT	
		or dried blood / body fluids on	Y	Ν	
L	the floor / top surfa			N	
^	•				
2.	Wash basin with ru	nning water or water stored in a	n Y	Ν	
2.	Wash basin with ru bucket with mug;	-		N	
	Wash basin with ru bucket with mug; soap, clean cloth or	towel available			
2. 3.	Wash basin with ru bucket with mug; soap, clean cloth or	towel available our coded bins are arranged in	Y Y	N N	

4.	Comments			
	Check actual practices at least once during the shift (Make a tick mark if it is done)			
5.	Hand-hygiene practiced	Y	Ν	
6.	Personal protective equipment are used correctly	Y	Ν	
7.	Needles /syringes are destroyed using needle hub cutter, needles collected in puncture proof container	Y	Ν	
8.	Waste segregated at source in correct colour coded bins	Y	N	
9.	All instruments are decontaminated, cleaned and sterilised. Chemical indicator is used	Y	N	
10.	Linen that is blood stained is decontaminated in a plastic bin. All linen are collected once a day by Class D worker and washed or sent to an external dhobi.	Y	N	
11.	All waste (infectious) waste is collected and put in appropriate pit (sharps in sharps pit/ infectious waste in respective pit) or collected by external agency	Y	N	
	Comments			

Appendix 6: Form 4: Record review (For Government and Private)

- 1. Name of the Facility:
- 2. Location
- 3. Date of assessment:
- 4. Name of the investigator:
- 5. Type of facility:
 - A. Government
 - i. UCHC
 - ii. UPHC
 - iii. ESI Dispensaries
 - iv. Health Kiosks

B. Private

- i. Private/Corporate Allopathy Hospital
- ii. Private/Corporate Allopathy Clinic
- iii. Private/Corporate AYUSH Hospital
- iv. Private/Corporate AYUSH Clinic
- v. NGO/FBO Allopathy Hospital
- vi. NGO/FBO Allopathy clinic
- vii. RMP
- viii. Other (specify)

5.A.1	Outcomes	March	Feb	Jan	Not
		2020	2020	2020	Applicable
1.	No registered in OPD in the morning				
2.	No registered in OPD in the evening				
	time				
3.	No. of ANC conducted				
4.	No. of outreach session conducted				
5.	No of emergencies				
6.	No of children with Acute malnutrition				
	referred to NRCs				
7.	No of children treated				

□ Diarrhea □ 8. Vital parameters (<i>if applicable</i>) No of births No of maternal deaths in catchment area □ No of newborn deaths in the past one year □ 9. No of children immunized □ 10. Pentavalent vaccine □ 10. Potta virus □ 10. No of TB patients on DOTS □ 10. No of TB patients on DOTS completing their treatment □ 11. No of patient attended OPD for any □ NCD □ □ □ □ □ □ □ □ □ □ □ 11. No of patient attended OPD for any □ NCD □ <t< th=""><th></th><th>□ Anemia</th><th></th><th></th></t<>		□ Anemia		
No of births Image: Second		Diarrhea		
No of maternal deaths in catchment area Image: Constraint of the past one year Image: Constraint of the past one year 9. No of children immunized Image: Constraint of the past one year Image: Constraint of the past one year 9. No of children immunized Image: Constraint of the past one year Image: Constraint of the past one year 9. No of children immunized Image: Constraint of the past one year Image: Constraint of the past one year Image: Constraint of the past one past of the past one in the past one past of the past one year Image: Constraint one year Image: Constrai	8.	Vital parameters (<i>if applicable</i>)		
area Image: second		No of births		
No of newborn deaths in the past one year		No of maternal deaths in catchment		
year Image: second		area		
9. No of children immunized - BCG vaccine - DPT vaccine - Pentavalent vaccine - Pentavalent vaccine - Hepatitis B vaccine - OPV - IPV - Rota virus - DT - HiB vaccine - MMR/MR - Measles 10. No of TB patients on DOTS No of TB patients on DOTS completing their treatment Image: Completing their treatment 11. No of patient attended OPD for any NCD • Diabetes- new • Hypertension - new • Diabetes- old/repeat • Total Diabetes		No of newborn deaths in the past one		
- BCG vaccine - DPT vaccine - Pentavalent vaccine - Hepatitis B vaccine - OPV - IPV - Rota virus - DT - HiB vaccine - MMR/MR - Measles 10. No of TB patients on DOTS No of TB patients on DOTS completing their treatment Image: Completing their treatment 11. No of patient attended OPD for any NCD • Diabetes- new • Hypertension - new • Diabetes-old/repeat • Total Diabetes		year		
 DPT vaccine Pentavalent vaccine Hepatitis B vaccine OPV IPV Rota virus DT HiB vaccine MMR/MR Measles 10. No of TB patients on DOTS 10. No of TB patients on DOTS completing their treatment 11. No of patient attended OPD for any NCD Diabetes- new Hypertension - new Diabetesold/repeat Hypertensionold/repeat Total Diabetes 	9.	No of children immunized		
- Pentavalent vaccine - Hepatitis B vaccine - OPV - IPV - Rota virus - DT - HiB vaccine - MMR/MR - Measles 10. No of TB patients on DOTS No of TB patients on DOTS completing their treatment		- BCG vaccine		
- Hepatitis B vaccine - OPV - IPV - Rota virus - DT - HiB vaccine - MMR/MR - Measles 10. No of TB patients on DOTS No of TB patients on DOTS completing their treatment		- DPT vaccine		
 OPV IPV Rota virus DT HiB vaccine MMR/MR Measles 10. No of TB patients on DOTS No of TB patients on DOTS completing their treatment 11. No of patient attended OPD for any NCD Diabetes- new Hypertension - new Diabetes-old/repeat Hypertension-old/repeat Total Diabetes 		- Pentavalent vaccine		
- IPV - Rota virus - DT - HiB vaccine - MMR/MR - Measles 10. No of TB patients on DOTS No of TB patients on DOTS completing their treatment Image: Completing their treatment 11. No of patient attended OPD for any NCD • Diabetes- new • Hypertension - new • Diabetes-old/repeat • Total Diabetes		- Hepatitis B vaccine		
 Rota virus DT HiB vaccine MMR/MR Measles 10. No of TB patients on DOTS No of TB patients on DOTS completing their treatment 11. No of patient attended OPD for any NCD Diabetes- new Hypertension - new Diabetes-old/repeat Hypertension-old/repeat Total Diabetes 		- OPV		
- DT - HiB vaccine - MMR/MR - Measles 10. No of TB patients on DOTS Image: Completing their treatment 11. No of patient attended OPD for any NCD Image: Complete set of the set of th		- IPV		
- HiB vaccine - MMR/MR - Measles 10. No of TB patients on DOTS No of TB patients on DOTS completing their treatment 11. No of patient attended OPD for any NCD • Diabetes- new • Hypertension - new • Diabetes-old/repeat • Total Diabetes		- Rota virus		
- MMR/MR - Measles - Measles 10. No of TB patients on DOTS - - No of TB patients on DOTS completing their treatment - - - 11. No of patient attended OPD for any NCD - - - • Diabetes- new - - - - • Diabetes- new - - - - • Diabetes- new - - - - - • Diabetes- old/repeat -		- DT		
- Measles - Measles 10. No of TB patients on DOTS No of TB patients on DOTS completing their treatment		- HiB vaccine		
10. No of TB patients on DOTS Image: Completing their treatment 11. No of patient attended OPD for any NCD Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment Image: Completing their treatment <th></th> <th>- MMR/MR</th> <th></th> <th></th>		- MMR/MR		
No of TB patients on DOTS completing their treatment Image: Completing their treatment 11. No of patient attended OPD for any NCD • Diabetes- new • Diabetes- new • Hypertension - new • Diabetes-old/repeat • Total Diabetes		- Measles		
their treatment Image: Constraint of the interval of the interva	10.	No of TB patients on DOTS		
11. No of patient attended OPD for any NCD • Diabetes- new • Hypertension - new • Diabetes-old/repeat • Hypertension-old/repeat • Total Diabetes		No of TB patients on DOTS completing		
NCD • Diabetes- new • Hypertension - new • Diabetes-old/repeat • Hypertension-old/repeat • Total Diabetes		their treatment		
 Diabetes- new Hypertension - new Diabetes-old/repeat Hypertension-old/repeat Total Diabetes 	11.	No of patient attended OPD for any		
 Hypertension - new Diabetes-old/repeat Hypertension-old/repeat Total Diabetes 		NCD		
 Diabetes-old/repeat Hypertension-old/repeat Total Diabetes 		• Diabetes- new		
Hypertension–old/repeatTotal Diabetes		• Hypertension - new		
Total Diabetes		Diabetes-old/repeat		
		• Hypertension-old/repeat		
Total Hypertension		Total Diabetes		
		Total Hypertension		
12. Records maintained about facility	12.	Records maintained about facility	I	
performance?		performance?		

Death register Image: No Applicable Emergency register MLC register Immunisation register Immunisation register Others specify Immunisation register Immunisation register Immunisation register Others specify Immunisation register Immunisation register Immunisation register Immunisation register 13. OPD prescription is given to the patient Immunisation Y Immunisation 14. Whether Patients have record of treatment? Y Not 5.A.2 Information Technology Immunisation registration Y Not 1. Unique identification number is given to each patient Y Not - At time of registration Y Not - Ordering tests N Applicable - At Pharmacy Y Not - Referring to other facilities Y Not - Births Y Not - Deaths Y Not - Computer Available/ Available/ - Online support Imagee <th></th> <th>Birth register</th> <th></th> <th>Yes</th> <th></th> <th>Not</th>		Birth register		Yes		Not
MLC register Immunisation register Communicable disease register Immunisation register Others specify Immunisation register 13. OPD prescription is given to the patient Y IA. Whether Patients have record of Y Immunisation Technology N Information Technology N Information Technology At time of registration Y At time of registration N Applicable - At time of registration N Applicable - At Pharmacy N Applicable - Follow-up N Applicable 2. What reports are sent to the Government? Not Births Y Not Deaths N Applicable - Communicable diseases Available/ Not 3. Equipments for data entry available? Available/ Not - Online support Available/ Not functional available		Death register		No		Applicable
Immunisation register Communicable disease register Immunisation register Others specify Immunisation register Immunisation register 13. OPD prescription is given to the patient Immunisation register Immunisation register 14. Whether Patients have record of treatment? Y Immunisation register 5.A.2 Information Technology Immunisation registration Y Not 1. Unique identification number is given to each patient N Applicable Applicable Immunisation Technology Immunisation registration Y Not Applicable 1. Unique identification number is given to each patient N Applicable Applicable 1. Unique identification number is given to each patient N Not Applicable 2. What reports are sent to the Government? Not Not Applicable 2. What reports are sent to the Government? Not Not Applicable 3. Equipments for data entry available? Available/ Not Available/ available 4. Passive data entry Immunicable data entry Immunic		Emergency register				
Communicable disease register Image: constraint of the specify Image: constraint of the specify 13. OPD prescription is given to the patient Image: constraint of the specify Image: constraint of the specify 14. Whether Patients have record of treatment? Image: constraint of the specify Image: constraint of the specify 5.A.2 Information Technology Image: constraint of the specify Image: constraint of the specify 1. Unique identification number is given to each patient Image: constraint of the specify Image: constraint of the specify Image: constraint of the specify 1. Unique identification number is given to each patient Image: constraint of the specify Image: constraint of the specify Image: constraint of the specify 1. Unique identification number is given to each patient Image: constraint of the specify Image: constraint of the spe		MLC register				
Others specify Others specify Image: specify Image: specify 13. OPD prescription is given to the patient Image: specify Image: specify 14. Whether Patients have record of treatment? Image: specify Image: specify 5.A.2 Information Technology Image: specify Image: specify 1. Unique identification number is given to each patient Image: specify Image: specify 1. Unique identification on umber is given to each patient Image: specify Image: specify 1. Unique identification on umber is given to each patient Image: specify Image: specify 1. Ordering tests Image: specify Image: specify Image: specify 1. Ordering tests Image: specify Image: specify Image: specify 2. What reports are sent to the Government? Image: specify Image: specify Image: specify 2. What reports are sent to the Government? Image: specify Image: specify Image: specify 3. Equipments for data entry available? Image: specify Image: specify Image: specify <th></th> <th>Immunisation register</th> <th></th> <th></th> <th></th> <th></th>		Immunisation register				
13. OPD prescription is given to the patient Image: space of the patient is given to the patient N 14. Whether Patients have record of treatment? Y N 5.A.2 Information Technology N N 1. Unique identification number is given to each patient Y Not - At time of registration Y Not - Ordering tests N Applicable - At Pharmacy N Not 2. What reports are sent to the Government? Y Not Births Paeaths Y Not 3. Equipments for data entry available? - Available/ Not - Online support Implicable - Available/ Not 4. Passive data entry - - Available / Not functional		Communicable disease register				
14. Whether Patients have record of treatment? N 14. Whether Patients have record of treatment? Y 5.A.2 Information Technology N 1. Unique identification number is given to each patient N - At time of registration Ordering tests N - At time of registration N Applicable - At Pharmacy N Applicable - Follow-up Not Applicable 2. What reports are sent to the Government? Not Births Y Not Deaths Not Applicable - Communicable diseases Not Applicable 3. Equipments for data entry available? Available/ Not - Online support Available / Not functional available 4. Passive data entry V Not functional Available /		Others specify				
14. Whether Patients have record of treatment? Image: constraint of treatment? N 5.A.2 Information Technology N N 1. Unique identification number is given to each patient Y Not - At time of registration Y Not - Ordering tests N Applicable - At Pharmacy N Applicable - Follow-up Y Not 2. What reports are sent to the Government? Y Not Births Y Not Applicable - Communicable diseases Y Not Applicable 3. Equipments for data entry available? Available/ Not - Computer Available/ Available/ Not - Online support Available / Not functional available	13.	OPD prescription is given to the patient		Y		
ireatment? N Impaire 5.A.2 Information Technology Impaire Impaire<				Ν		
5.A.2 Information Technology Information Technology 1. Unique identification number is given to each patient Y Not - At time of registration Y Not - Ordering tests N Applicable - At Pharmacy N Applicable - Referring to other facilities N Applicable - Follow-up Not Applicable 2. What reports are sent to the Government? Y Not Births Y Not Applicable - Deaths N Applicable - Communicable diseases Not Applicable - Computer Available/ Not - Online support Functional available - Not functional Available / Not functional - Passive data entry Unique identification Unique identification	14.	Whether Patients have record of		Y		
1. Unique identification number is given to each patient Patient Not - At time of registration PY Not - Ordering tests N Applicable - At Pharmacy Not Applicable - Referring to other facilities Not Applicable - Follow-up Not Applicable 2. What reports are sent to the Government? Not Applicable - Births Y Not - Deaths Not Applicable - Communicable diseases Not Applicable - Computer Available? available - Online support Functional available - Not functional Available / Not functional 4. Passive data entry I I I I		treatment?		Ν		
patient - At time of registration - Y Not - Ordering tests - N Applicable - At Pharmacy - Referring to other facilities - Follow-up - Follow-up - Follow-up - Y 2. What reports are sent to the Government? - Y Births - Y Not - Deaths - Y Applicable - Communicable diseases - Y Applicable 3. Equipments for data entry available? - Available/ Not - Computer - Computer - Available/ available - Not functional - Available - Available - Available	5.A.2	Information Technology				
patient - At time of registration - Y Not - Ordering tests - N Applicable - At Pharmacy - Referring to other facilities - Follow-up - Follow-up - Follow-up - Y 2. What reports are sent to the Government? - Y Births - Y Not - Deaths - Y Applicable - Communicable diseases - Y Applicable 3. Equipments for data entry available? - Available/ Not - Computer - Computer - Available/ available - Not functional - Available - Available - Available			T			
At time of registrationImage: Y model of the second of the se	1.		ach			
 Ordering tests At Pharmacy Referring to other facilities Follow-up What reports are sent to the Government? Births Deaths Communicable diseases Equipments for data entry available? Computer Online support Available/ Not Available/ Not Available / Not functional Passive data entry 						
 At Pharmacy Referring to other facilities Follow-up Follow-up What reports are sent to the Government?		- At time of registration			Y	Not
 Referring to other facilities Follow-up What reports are sent to the Government? Births Deaths Communicable diseases Computer Computer Available/ Not Available / Not Available / Not functional Available / Not functional Passive data entry 		- Ordering tests			Ν	Applicable
- Follow-up- Follow-up- Second		- At Pharmacy				
2. What reports are sent to the Government? Image: Sent to the Government? Image: Sent to the Government? Image: Births Image: Births Image: Sent to the Government? Image: Sent to the Government? Image: Not Applicable Image: Deaths Image: Deaths Image: Sent to the Government? Image: Sent to the Government? Image: Not Applicable Image: Deaths Image: Communicable diseases Image: Sent to the Government? Image: Sent to the Government? Image: Applicable 3. Equipments for data entry available? Image: Sent to the Government? 3. Equipments for data entry available? Image: Sent to the Government? 3. Equipments for data entry available? Image: Sent to the Government? Image: Sent to the Government Image: Sent to to to to to to to t		- Referring to other facilities				
Image: Sector of the sector		- Follow-up				
□Births□NApplicable□Deaths□NApplicable□Communicable diseases□3.Equipments for data entry available?-Available/Not-Computer□Available/Not-Online support-FunctionalavailableNot functional-Not functional4.Passive data entry	2.	What reports are sent to the Government?				
Deaths Deaths Communicable diseases Image: Communicable diseases 3. Equipments for data entry available? - Computer - Computer - Online support Image: Computer Functional - Available / Not Not functional 4. Passive data entry		□ Births			Y	Not
Communicable diseases Image: Communicable diseases 3. Equipments for data entry available? - Computer - Online support - Online support - Available/ - Available / - Not - Not functional - Available / - Not functional		Deaths			Ν	Applicable
3. Equipments for data entry available? Image: Available / Not available - Computer Image: Available / Not available - Online support Functional Image: Available / Not functional Not functional 4. Passive data entry Image: Available / Not functional						
- Computer - Available/ Not - Online support Functional available - Available / Not functional - Not functional 4. Passive data entry						
 Online support Functional Available / Not functional 4. Passive data entry 	3.					
4. Passive data entry		_				
4. Passive data entry		- Online support			Functional	available
4. Passive data entry					Available /	
					Not functional	
- Aggregate numbers only \Box Y Not	4.	Passive data entry				
		- Aggregate numbers only			Y	Not
- Facility data 🗆 N Applicable		- Facility data			Ν	Applicable
- Community data		- Community data				

Appendix 7: House-listing and Screening

Wa	ard CEB Interviewer Household							
	Section 1 – House-listing & Geolocation							
1)	Login using your credentials (fieldworker name)							
	Select urban ward name.							
3)	Select CEB Block name							
4)	Date of Interview (dd/mm/yyyy)							
5)	Individual house/ apartment							
6)	No of households within house or within apartment (based on operational kitchen)							
7)	Household sequence # (Auto generate)							
	Household Availability.							
	Options : $1 =$ Primary Respondent Available							
	2 = Door locked-if so time to visit							
	3 = Primary Respondent is not available – if so time to visit							
	4 = Busy -if so time to visit							
	5 = Refusal (They are not interested to take survey)							
	6 = Others/Remarks (Specify)							
9)	Household address (Door #/Street address /pincode & landmark):							
-								
	Primary & Secondary mobile:							
	Head of the family: Choice of language for interview:							
	Primary respondent:							
10)	Friend/neighour 1 Household details (Name, Door #/Street address /pincode & landmark, Primary mobile) :							

S1.	Name	Relation	Sex	Age	D.O.B.	If fem	If fem and		If age>30
No		ship		(completed	(if age	and aged	aged 18-	age <60	yrs, has
		to head		yrs)	<5 yrs)	15-49,	52,	months, is	DM or HT
						currentl	pregnancy	currently	– yes /no
						У	in the last	sick or	
						pregnant	3 years –	was sick	
						-	yes / no?	in the last	
						yes/no?		1 month-	
						-		yes / no?	
1		Head							
2									
3									
4									
5									

Relationship to head, Options: 1 = Head, 2 = Spouse, 3 = Son/Daughter, 4 = Parent, 5 = Parent-in-law, 6 = Brother/Sister, 7 = Brother/Sister-in-law, 8 = Grandson/Granddaughter, 9 = Domestic Help, 10 = Other Relatives

- 1. Generate unique ID for the household:
- 2. Save and Submit form.
- **3.** Creation of Dashboard to review survey details till date based on household sequence # /street address/ area/ date of interview.

Appendix 8: Every 7th household survey

Section 3 - Household	Section 3 - Household Characteristics							
	Date of interview _	(dd/mm/yyyy)						
3.1 Name of head of he	ousehold:							
3.2 Contact numbers:	Landline							
	Mobile-1	Mobile-2						
3.3 Choice of language	for interview Kannada-1 English- 2 Urd	u - 3 Others $- 4$;						

3.4 If you out-migrate, friend/neighbour we may contact (name/address/contact no)

3.5 Type of house (on observation)	1. Pucca, 2. Semi Pucca, 3. Kachha
3.6 Availability of drinking water source	1. Within the premises 2. Near the premises 3. Away 4. Others (Specify)
3.7 Ownership status of house	1. Owned, 2. Rented/leased, 3. Other (Specify)
3.8 Solid waste segregated at source	1. Yes 2. No
3.9 Refrigerator	1. Yes 2. No
3.10 Computer/Laptop	1.Yes: With internet 2. Without internet 3. No
3.11 Motorized wheelers	1. Four-wheeler 2. Three-wheeler 3. Two-wheeler
3.12 Total number of mobile/phones – smartphones and basic phones separately	Basic Phones Mobile Phones — — — —
3.13 Toilet facility	1. Improved not shared 2. Shared facility 3. Unimproved 4. No facility/uses open space
3.14 Cooking Fuel	 LPG/electricity 2. Kerosene 3. Solid (wood,etc) Other
3.15 Religion	1. Hindu2. Muslim3. Christian4. Other (specify)
3.16 Caste	1. Scheduled caste2. Scheduled Tribe3. Other backward caste4. other
3.17 May I please see the BPL card of your family	1. Seen Yes 2. No

3.18 Where do you/your family members usually go for health care?	1. Government, Specify 2. Private, Specify

Section 4: Household listing (Listing of household member (usual residents only))

4.1. Sl. No	4.2. Name	4.3. Relationshi p to head	4.4 Sex	4.5.Age (complet ed yrs)	4.6.D.O. B. (if age <5 yrs)	4.7. Marital Status	4.8.Educat ion	4.9.Occupation
1		Head						
2								
3								
4								
5								
6								
7								

Sl N o	Name	4.10.Income per month	4.11. Health Insurance (see AB-ArK card)	4.12. Did you use any of the health insurance in the past 1 year for any illness? 1-Yes 2-No	much did you spend towards illness?: 4.13b. how much was insurance	Illness in the last 2
1						1 2
2						1 2
3						1 2
4						1 2
5						1 2
6						
7						

Codes: **Sex:** 1. Male 2. Female 3. Other; **Marital Status:** 1. Currently married; 2. Widow/ widower/ divorced/ separated / deserted; 3. Never married; 4. N/A; **Education:** 1. Illiterate, 2. Primary, 3. Middle, 4. High School, 5. PUC, 6. Graduate/Diploma, 7. Postgraduate 8. N/A; **Occupation:** 1. Government, 2. Private, 3. Self-employed, 4. Business, 5. Daily wages, 6. Retired, 7. Student, 8. Home maker

Insurance: 1. Ayushman Bharat/Arogya Karnataka, 2. Employees state Insurance scheme (ESI)/CGHS, 3. Other privately purchased health insurance, 4. Medical reimbursement from employer, 5. Others.

	Options	Last Immunization
5.1 Where did you go for?	1 – Government, Specify 2 – Private. Specify	
5.2 What was the primary reason for preference?	 Close by Less cost Free of cost Trust/Good doctor Timely service All facility at one place Others-Specify 	
5.3 How much time did you have to spend during the clinic visit?	 < 30 minute 30 minutes - 1 hour 1 hour - 2 hours 2 hours - 4 hours > 4 hours 	
5.4 Overall, were you satisfied with child health services provided?	Please show me on this visual scale of 0 to 10	
5.5 Child hospitalized?	1. Yes 2. No	

Section 5 - Child Immunization (under 24 months)

Section 6 - Lifestyle (>18 years)

6.1 Do you currently smoke any tobacco products, such as	1. Yes	
cigarettes, cigars, pipes, beedis, hookahs?	2. No	
6.1a. If in case having smoking habits, how many days in a	1. Daily	
week do you smoke?	2. 5–6 days per week	
	3. 1–4 days per week	
	4. 1–3 days per month	
	5. Less than once a	
	month	
6.2 Do you currently use any smokeless tobacco (chewing	1. Yes	
tobacco)	2.No	
6.2a. If Yes, how many days in a week do you consume	1. Daily	
	2. 5–6 days per week	
	3. 1–4 days per week	
	4. 1–3 days per month	
	5. Less than once a	
	month	
	1. Yes	
6.3. Do you consume alcoholic drink	2. No	
	1. Daily	
	2. 5–6 days per week	
	3. 1–4 days per week	
	4. 1–3 days per month	
6.3a. If in case alcohol drinking habit. How many days in a	5. Less than once a	
week do you drink?	month	
6.4. In a typical week, on how many days do you eat fruit?	days	
6.5. How many servings of fruits do you eat on one of those	1. Lessthan 100 gm	5.
days?	2. $100 \text{gm} - 200 \text{gm}$	5.
uuys.	3. $200 \text{gm} - 300 \text{gm}$	
	4. Greater than	
	300gm	
6.6. In a typical week, on how many days do you eat	days	
vegetables?	auj 5	
6.7. How many servings of vegetables do you eat on one of	1. Lessthan 100 gm	
those days?	2. 100gm – 200gm	
	3. 200gm – 300gm	
	4. Greater than	
	300gm	
6.8. How often do you add salt to your food before you eat	1. Always	
it or as you are eating it?	2. Often	
	3. Sometimes	
	4. Rarely	
	5. Never	
	6. At food	
	preparation	
	7. Don't know	
	7. DOINT KNOW	

6.9. Are you a member of any voluntary organization?	 Mahila Arogya Samiti Self-Help Group Other (specify) Never
6.10.Do you go for brisk walk?	1. Yes 2. No
6.11.In a week, how many days do you go for a brisk walk?	days
6.12. What is the distance to the nearest public park for walking/exercise(kms)	(Kms)
6.13. Have you gone for an eye check-up on your own?	1. Yes 2. No

Section 7 – Two-week morbidity details

7.1.Acute Ailments	7.2.Name of ailment	7.3.Did you seek any treatment? 1. Yes 2. No	7.4.If no, what was the reason?	7.5.Where did you avail treatment ? 1. Govt 2. Private 3. None	7.6.Which system of medicine?1. Allopathy2. Ayurveda3. Homeo4. Others	7.7.Name of the facility and location	7.8.Did you change from usual place of care to other place for this ailment	7.9.Reason s for changing place of care	7.10. Were you satisfied with the services?1. Yes2. No	7.11.Reasons for satisfaction/di ssatisfaction with service
Ailment 1 Place 1 Place 2 Place 3 Place 4										
Ailment 2 Place 1 Place 2 Place 3 Place 4										
Ailment 3 Place 1 Place 2 Place 3 Place 4										

7.4. If no, what was the reason? 7.9. Reason-changing place of care

1. Home Remedies

- 2. OTC (over the counter)
- 3. Medicine available at home

4. Others (Specify)

- - 1. Far away
 - 2. High Cost
 - 3. Distrust Doctor
 - 4. Slow/late service
 - 5. Less facilities
 - 6. Others (Specify)

7.11 Reason for satisfaction

- 1. Close by
- 2. Less cost
- 3. Free of cost
- 4. Trust/Good Doctor
- 5. Timely Service
- 6. All facility in one place
- 7. Others (specify)

7.11. Reason for dissatisfaction

- 1. Far away
- 2. High Cost
- 3. Distrust Doctor
- 4. Slow/late service
- 5. Less facilities
- 6. Others (Specify)

Acute Ailments	7.12.Amou nt spent on consultatio n/payment to doctor if Govt.	7.13.W as any investig ation done? 1. Yes 2. No	7.14.Am ount spent on tests/ investigat ions (enter zero (0) if no expenses)	7.15.Were youprescribedmedicines?1. Yes2. No	7.16. Amou nt spent on drugs	7.17.Cu mulativ e amount spent if breakdo wn not known	7.18. Amou nt Spent on Travel	7.19.Mod e of travel 1-Bus 2-Car 3-2 wheeler 4-Walk 5-others	7.20.Dis tance to health facility (km)	7.21 Time taken to reach the facility	7.22.Time of consultati on (7.23.Time taken to consult the doctor (in minutes)	 7.24. how do you manage your routine medical expenses? 1. income 2. savings 3. borrowed 4. Selling Property 5. Selling Jewellery 6. Insurance
Ailment 1 Place 1 Place 2 Place 3 Place 4													
Ailment 2 Place 1 Place 2 Place 3 Place 4 Ailment 3													

Formative Research for CPHC in Mysuru City

-	Place							
	1							
-	Place							
	2							
-	Place							
	3							
-	Place							
	4							

7.21 Options:

7.22 Options :

7.23 Options :

3. 1 hour - 2 hour

- 1. less than 30 minutes
- 2. 30 minutes 1 hours
- 3. 1 hour 2 hour
- 4. 2-hour -4 hour
- 5. Greater than 4 hours

- 1. Morning
- 2. After noon
- 3. Evening
- 4. Night

4. 2-hour -4 hour

1. less than 30 minutes

2. 30 minutes - 1 hours

5. Greater than 4 hours

	Options		Delivery	PNC
If within Mysore city, complete section 8		ANC		
8.1. Which type of health facility did you use regularly	1. Government, Specify			
for maternal health care during your most recent	2. Private Specify			
pregnancy/Abortion?	3. Traditional birth attendant			
	4. Other, Specify			
8.2. What was the primary reason for preference?	1. Close by			
	2. Less cost			
	3. Free of cost			
	4. Trust/good doctor			
	5. Timely service			
	6. All facilities under one roof			
	7. Other			
8.3. Did you receive any health record/ Thayi card?	1. Yes			
	2. No			
8.4. How long did it take you to reach the facility?	minutes			
8.5. What was the mode of transport to the facility?	1. Bus			
	2. Car			
	3. 2- Wheeler			
	4. By walk			
	5. Other			
8.6. Did you miss any scheduled visits?	1. Yes			
	2. No			-
8.7. Were any complications detected in mother?	1. Yes			
	2. No			
8.8. Were you referred to another hospital for treatment	1. Govt			
of these complications?	2. Private			
	3. None			
8.9. Were any complications detected in baby?	1. Yes			
	2. No			
8.10. Were you referred to another hospital for	1. Govt			
treatment of these complications?	2. Private			
	3. None			
8.11. What was type of hospital that you went to?	1. Govt			
	2. Private			
	3. None			
8.12. Type of delivery	1. Normal			
	2. C-section			
	3. Other			
8.13. In total, how much did your household spend for maternal health services during your last pregnancy?				
8.14. Overall, how satisfied were you with the maternal	Please show me on this			
health services you received?	visual scale of 0 to 10			
8.15. Did you receive direct cash transfer? (JSY/Other)	1. Yes			
	2. No			
	2.110			

Section 8: Maternal Health (special survey) (Place of delivery – Within Mysore City / Outside Mysore)

8.16. How do you manage your routine medical expenses?	1.Income2.Savings3.Borrow from family/other4. Selling Property5. Selling Jewellery6. Insurance
--	--

Section 9 - Child Health (special survey)

	Options	Morbidity in last 1 month = Yes			
		ARI	ADD	Other illness	
		(Yes / No)	(Yes / No)	(Yes / No)	
9.1. Where did you go for?	1. Government, specify				
	2 Driverte Streetfor				
0.2. What was the mimory reason	2. Private, Specify				
9.2. What was the primary reason	 Close by Less cost 				
for preference?	2. Less cost 3. Free of cost				
	4. Trust/Good doctor				
	5. Timely service				
	6. All facility at one place				
	7. Other				
9.3. How much time did you have	1. less than 30 minutes				
to spend during the clinic visit?	2. 30 minutes - 1 hours				
	3. 1 hour - 2 hour				
	4. 2-hour -4 hour				
	5. Greater than 4 hours				
9.4. Child hospitalized?	1. Yes				
	2. No				
9.5. If Yes, place of hospitalization					
Name /Location					
9.6.In total, how much did your					
household spend for child health services?					
IP					
IT OP					
9.7. Overall, are you satisfied with	Please show me on this				
child health services, provided?	visual scale of 0 to 10				
9.8. How do you manage your	1. Income				
routine medical expenses?	2. Savings				
L. L	3. Borrow from				
	family/other				
	4. Selling Property				
	5. Selling Jewellery				
	6. Insurance				

		Diabetes	Hypertension
10.1.Where do you routinely go for your doctor consultation?	 Government, Specify Private, Specify Others 		
10.2. Which system of medicine?	 Allopathy Ayurveda Homeopathy Others 		
10.3. If government, name of facility?			
10.4. Location			
10.5. Reason for preference	 Close by Less cost Free of cost Trust/Good doctor Timely service All facility at one place Other 		
10.6. If private, name of facility?			
/Doctor name/Location 10.7. Location			
10.8. Reason for preference	 Close by Less cost Free of cost Trust/Good doctor Timely service All facility at one place Other 		
10.9. Routinely where do you get your fasting blood sugar test done? (only for DM)	1. Govt 2. Private		
DM) 10.10. Reason for preference	3. Others		
10.11. Where did you go for last check up?			
10.12. In the last 6 months, how many doctors have you consulted?			
10.13. Do you take regularly treatment for this condition?	1. Yes 2. No		
10.14. In the last 7 days, on how many days have you missed taking medicines?			
10.15. Routinely where do you get your medicines from?	1. Govt 2. Private 3. None		

Section 10 - Non-communicable Diseases (in the past 1 year) (special survey)

10.16. Reason for preference for buying medicines 10.17. Have you ever got your eye check-up done	 1.Close by 2.Less cost 3.Free of cost 4.All medicines are available 5. Other(Specify) 1. Yes 2. No 		
10.18. Have you got blood check done	1. Yes		
for kidney disease10.19. Have you got blood check donefor cholesterol	2. No 1. Yes 2. No		
10.20. Have you ever got ECG done	1. Yes 2. No		
10.21. Has your doctor ever told you that you have any complications?	1. Yes 2. No		
10.22. If yes, where were you referred to?	 Government Private 		
10.23. Where did you go? Name and location	 Government (Specify) Private (Specify) 		
10.24 Please tell me on a scale of 0 to 10 how satisfied are you with govt hospital , where 0 means not satisfied at all and 10 means you are completely satisfied	$\begin{array}{cccccccccccccccccccccccccccccccccccc$		
10.25. Reason for the above score			
10.26. Please tell me on a scale of 0 to 10 how satisfied are you with Private hospital , where 0 means not satisfied at all and 10 means you are completely satisfied	$\begin{array}{cccccccccccccccccccccccccccccccccccc$		
10.27. Reason for the above score			
10.28. Most recent levels		FBS PPBS RBS HbA1c	SBP- DBP –
10.29 How do you manage your routine medical expenses?	 Income Savings Borrow from family/other Selling Property Selling Jewellery Insurance 		

- 10.30. Have you had any medical check-up for mouth cancer? (for >18 years) 1- Yes, 2- No
- 10.31. Have you had any medical check-up for cervical cancer? (for >30 years, women) 1 -Yes, 2-No
- 10.32. Have you had any medical check-up for breast examination/mammography/other test) breast cancer? (for >18 years, women) 1 -Yes, 2 -No

10.33.Record Blood Pressure Measured: SBP1:_____ DBP1:_____ Pulse1_____

> SBP2:_____ DBP2:_____ Pulse2_____